

ÜLLE VÕHMA

Association between personality traits,
clinical characteristics and pharmacological
treatment response in panic disorder



ÜLLE VÕHMA

Association between personality traits,
clinical characteristics and pharmacological
treatment response in panic disorder



Department of Psychiatry, Institute of Clinical Medicine, University of Tartu, Tartu, Estonia

The dissertation is accepted for the commencement of the degree of Doctor of Philosophy (medicine) on June 25, 2019 by the Council of the Faculty of Medicine, University of Tartu.

Supervisors: Eduard Maron, MD, PhD, Professor,
Department of Psychiatry, Institute of Clinical Medicine, Faculty of
Medicine, University of Tartu, Tartu, Estonia

Anu Aluoja, PhD, Associate Professor,
Department of Psychiatry, Institute of Clinical Medicine, Faculty of
Medicine, University of Tartu, Tartu, Estonia

Innar Tõru, MD, PhD, Associate Professor,
Department of Psychiatry, Institute of Clinical Medicine, Faculty of
Medicine, University of Tartu, Tartu, Estonia

Reviewers: Jaanus Harro MD, PhD, Professor and Head,
Department of Neuropsychopharmacology, Institute of
Psychology, Faculty of Social Sciences, University of Tartu,
Tartu, Estonia

Kirsti Akkermann PhD, Associate Professor,
Department of Clinical Psychology, Institute of Psychology, Faculty
of Social Sciences, University of Tartu, Tartu, Estonia

Opponent: Virginija Adomaitiene, MD, PhD, Professor, Head of Psychiatry
Department, Lithuanian University of Health Sciences, Medical
Academy, Kaunas, Lithuania

Commencement: September 03, 2019

This study was supported by the grant from Estonian Science Foundation No. 4614; the targeted financing from Estonian Ministry of Education and Research, No SF0180125s08, Estonian Science Foundation grant 7034, and “Institutional Research Funding” project IUT20-45.

ISSN 1024-395X
ISBN 978-9949-03-122-1 (print)
ISBN 978-9949-03-123-8 (pdf)

Copyright: Ülle Võhma, 2019

University of Tartu Press
www.tyk.ee

CONTENTS

LIST OF ORIGINAL PUBLICATIONS	7
ABBREVIATIONS.....	8
1. INTRODUCTION	9
2. REVIEW OF THE LITERATURE.....	11
2.1. Clinical characteristics and course of panic disorder	11
2.2. Risk factors of panic disorder	13
2.3. Pharmacotherapy of panic disorder and treatment outcome	14
2.4. Personality and panic disorder	15
2.5. Personality and treatment outcome in panic disorder	18
3. RATIONALE FOR THE STUDIES.....	21
4. AIMS OF THE STUDIES	23
5. MATERIALS AND METHODS.....	24
5.1. Subjects.....	24
5.2. Ethical considerations	25
5.3. Measures	25
5.3.1. The Mini International Neuropsychiatric Interview (M.I.N.I.5.0.0.).....	25
5.3.2. Swedish universities Scales of Personality (SSP).....	26
5.3.3. The revised NEO Personality Inventory (NEO-PI-R).	26
5.3.4. Panic Disorder Severity Scale (PDSS)	27
5.3.5. The Clinical Global Impression scale (CGI)	27
5.3.6. Hamilton Anxiety Scale (HAM-A).....	28
5.3.7. Toronto Side-effect Scale (TSES)	28
5.3.8. The Emotional State Questionnaire (EST-Q)	28
5.3.9. Sociodemographic measures	28
5.4. Treatment	28
5.5. Statistical analysis.....	29
6. RESULTS AND DISCUSSION	31
6.1. Study I: Factor structure of the Swedish universities Scales of Personality and position within the five-factor model in an Estonian sample	31
6.1.1. The gender differences for the SSP	31
6.1.2. The principal component analysis of the SSP scales	32
6.1.3. The correlations between the SSP and NEO-PI-R scales	32
6.1.4. Discussion.....	33
6.2. Study II: Evaluation of personality traits in PD using SSP.....	36
6.2.1. Demographic and clinical characteristics	36
6.2.2. Personality differences between total PD group and healthy controls	37
6.2.3. Personality traits in patients with pure and comorbid PD.....	40

6.2.4. Associations of clinical and demographic factors with the personality	40
6.2.5. Correlation data	41
6.2.6. Discussion.....	41
6.3. Study III: Association between personality traits and escitalopram treatment efficacy in PD	44
6.3.1. Demographic and clinical characteristics	44
6.3.2. Changes in SSP sub-scale scores following escitalopram treatment in PD	45
6.3.3. Correlation data	46
6.3.4. Discussion.....	49
7. GENERAL DISSCUSSION	52
8. CONCLUSIONS.....	56
9. REFERENCES	57
SUMMARY IN ESTONIAN	70
ACKNOWLEDGEMENTS	77
PUBLICATIONS	78
CURRICULUM VITAE	105
ELULOOKIRJELDUS.....	106

LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following original publications, which are referred to in the text by Roman numerals I–III.

- I Aluoja A, Voogne H, Maron E, Gustavsson P, **Võhma Ü**, Shlik J. Personality traits measured by the Swedish universities Scales of Personality: Factor structure and position within the five-factor model in an Estonian sample. *Nordic J of Psychiatry* 2009; 63 (3): 231–236.
- II **Võhma Ü**, Aluoja A, Vasar V, Shlik J, Maron E. Evaluation of personality traits in panic disorder using Swedish universities Scales of Personality. *J Anx Disord* 2010; 24: 141–146.
- III **Võhma Ü**, Raag M, Tõru I, Aluoja A, Maron E. Association between personality traits and escitalopram treatment efficacy in panic disorder. *Nordic J of Psychiatry* 2017; 4: 1–8.

Author's contribution to the preparation of the original publications:

Publication I: Participation in setting the aims and designing the study; interpretation of results; writing the manuscript as a coauthor.

Publication II: Participation in setting the aims and hypotheses, designing the study; participation in collecting and analysing the data; interpretation of results; writing the manuscript as the main author.

Publication III: Participation in setting the aims and designing the study; organising the study and collecting the data; participation in analysis of the data; interpretation of results; writing the manuscript as the main author.

ABBREVIATIONS

APA	American Psychiatric Association
BPD	Bordeline Personality Disorder
CBT	Cognitive Behavioural Therapy
CCK	Cholecystokinin
CGI	Clinical Global Impression scale
CGI-I	Clinical Global Impression – Improvement scale
CGI-S	Clinical Global Impression – Severity scale
CI	Confidence Interval
CSF	Cerebrospinal Fluid
DSM-IV	Diagnostic and Statistical Manual of Mental Disorders(4th Edition)
ED	Emergency Department
EST-Q	Emotional State Questionnaire
HAM-A	Hamilton Anxiety Scalet
5-HTTLPR	the serotonin transporter gene-linked polymorphism
ICD-10	the International Statistical Classification of Diseases, 10th Revision
KSP	the Karolinska Scales of Personality
MINI	Mini-International Neuropsychiatric Interview
N	Neuroticism
NEO-PI-R	Revised Neuroticism-Extraversion-Openness Personality Inventory
OR	Odds ratio
PD	Panic disorder
PDSS	Panic Disorder Severity Scale
SNRI	Selective Serotonin-Norepinephrine Reuptake Inhibitor
SSP	the Swedish universities Scales of Personality
SSRI	Selective Serotonin Reuptake Inhibitors
SSTI	selektiivne serotoniini tagasihaarde inhibiitor
TCI	Temperament and Character Inventory
TSES	Toronto Side-effect Scale

1. INTRODUCTION

Panic disorder (PD) was officially introduced into the psychiatric nomenclature as a distinct illness in 1980 with the publication of the Diagnostic and Statistical Manual of Mental Disorders, third edition (APA, 1980). The central clinical features of PD are spontaneous panic attacks, involving numerous psychological and physical symptoms. PD is generally considered a chronic or intermittent psychiatric condition which is often disabling (Keller et al., 1994, Batelaan et al., 2010). The illness leads to an increased use of health care services (Rief et al., 2005) as well as a significant loss in the quality of life and psychosocial functioning (Keller et al., 1994, Kessler et al., 2006; Roy-Burne et al., 2006, Mendlowicz et al., 2000).

Genetic, biological, environmental and psychosocial factors, including personality traits were found to play an important role in the development of the PD and influence treatment response (Andrisano et al., 2012, Carrera et al., 2006). Furthermore, a study by Powers and Westen (2009) produced results that suggest the importance of personality in understanding the different phenotypes of panic patients (high functioning, emotionally dysregulated, inhibited/ avoidant, somatizing patients) and in making better predictions about responses to treatment (Powers, 2009). This point of interest was an important motivator for this dissertation.

The expression of personality traits mainly related to anxiety and mood disorders is proposedly connected to serotonin mechanisms (Takano et al. 2007). High levels of Neuroticism have been associated with increased serotonin binding within the thalamus (Takano et al., 2007) and 5HT_{1A} receptor (Hirvonen et al., 2015)

Personality traits can have some influence on the outcome of pharmacotherapy, which in turn can produce changes in personality. Studies have revealed contradictory results. Some initial reports on PD have indicated that a large improvement in panic symptoms after anxiolytic medication with alprazolam or diazepam is accompanied by significant changes in neuroticism-related personality traits (Reich et al., 1991, Noyes et al., 1986). On the contrary, studies by Carrera et al. (2006) postulated that none of the personality traits of panic disorder patients, assessed by the Neuroticism-Extraversion-Openness Five-factor Personality Inventory (NEO-FFI) were related either to clinical severity or to a short-term response to SSRI treatment (Carrera et al., 2006).

The relation between personality traits and the treatment outcomes of PD remains an important topic for physicians because, although progress has been made in the development of effective pharmacological and psychotherapeutic interventions, still not all patients respond to treatment (Katon et al., 2006). The high rate of treatment failure and relapses in treating PD patients possibly suggest personality diathesis (Powers and Westen, 2009). A more effective comprehension of the relationship between personality factors and PD can help to give a greater understanding of etiology, identify patients at risk of suffering

from more resistant symptoms, and help to both plan treatment and prevent these conditions.

The impact of personality traits on the risk of panic disorder and on its treatment, as well as the stability of a personality disorder diagnosis, has continually been an object of discussion (Johnson et al., 2006; Mennin et al., 2000).

2. REVIEW OF THE LITERATURE

2.1. Clinical characteristics and course of panic disorder

Panic disorder (PD) is a serious anxiety disorder characterised by recurrent panic attacks and associated fearful anticipation of panic and its consequences, and frequently developing agoraphobia (DSM-IV, ICD-10). Panic attacks are defined as rapidly escalating occurrences of at least four out of 13 distressing symptoms including palpitations, sweating, trembling or shaking, shortness of breath or smothering, feeling of choking, chest pain or discomfort, nausea or abdominal distress, feeling dizzy, unsteady, lightheaded, or faint, derealisation, depersonalisation, fear of losing control or going crazy, fear of dying, paresthesias and chills or hot flushes. The identification of PD requires careful differential diagnosis from panic-like symptoms and autonomic arousal resulting from the direct physiological effects of substance use or a general medical condition and from symptoms of anxiety better explained by another mental health disorder. On the other hand, patients with PD have an increased risk of other psychiatric as well as medical morbidity. There is a high degree of PD comorbidity with other mental health disorders: 30–63% for major depression, 29.5–58.2% for agoraphobia, 20–75% for social phobia, 20% for generalised anxiety disorders, 14% for obsessive-compulsive disorder, 13% for bipolar disorder and 6% for post-traumatic stress disorder (Goisman et al., 1994; Chen and Dilsaver, 1995; Pelissolo and Lepine 1998; Eaton et al., 1994; Wittchen et al., 1998; Weissman et al., 1997; Kessler et al., 1998; Kessler et al., 2006; Lamers et al., 2011; Preti et al., 2018). Substance abuse is also a common comorbid disorder in up to 36% of cases according to Epidemiological Catchment Area data (Regier et al., 1998). In addition, PD often occurs together with cardiac, gastrointestinal, respiratory and neurological disorders (Zaubler and Katon, 1998).

According to epidemiological surveys, sporadic panic attacks occur in the population at 7–15% the lifetime prevalence rate (Pelissolo and Lepine, 1998; Eaton et al., 1994), and are often associated with substantial morbidity even in the absence of a full clinical manifestation of PD (Klerman et al., 1991). The lifetime prevalence of PD rates are between 1.5% and 3.5%, while twelve-month prevalence rates in females were found to be higher than in males (Eaton et al., 1994). In Estonia, the prevalence of panic symptoms among primary care patients is different among men and women, accordingly 5.5% for men and 8% for women (King et al., 2008) – a tendency that has been demonstrated in several studies. The age of onset of PD is usually in the mid-twenties, with hazard rates for females ranging from 25 to 34 years and for males between 30 and 44 years (Wittchen and Essau, 1993). As with gender, data about PD is also different with age. Panic attacks were observed in 3% of youths aged from 9 to 17 years (Goodwin and Gotlib 2004) and PD was diagnosed in 10–13% of clinically referred children and adolescents (Doerfler et al., 2007, Masi et

al.,2000). Lower rates of panic disorder are typically seen in persons aged 65 and older (Robins, 1991, Kessler et al., 2006, Yates, 2009).

PD has been recognised as a chronic condition with a fluctuating course (Liebowitz 1997; Pollack and Otto, 1997). Some studies have addressed longitudinal aspects; for example, in a three-year follow-up only 10% of patients with PD were symptom-free (Noyes et al., 1990) and only 12% of PD patients were in full remission after five years (Faravelli et al., 1995). At follow-up, study revealed that 25–50% of PD patients relapse within 6 months after discontinuation of drug treatment (Pull and Damsa, 2008). There is evidence that the more important predicting factors of the course of panic disorder in the clinical population are not the severity and frequency of panic attacks, but rather long duration and agoraphobia at baseline (Katschnig and Amering, 1998). Predictors of remission were found to be female gender, the absence of ongoing difficulties, subthreshold panic and a low initial frequency of attacks (Katon, 1987, Kessler 2006).

PD can be a seriously disabling disorder that causes impairment in social, personal and occupational functioning and a significant loss of quality of life (Candilis et al., 1999). The rate of those PD patients who are dependent on welfare or disability benefits exceeds 25% in some studies (Katerndahl and Realini, 1997). Furthermore, medical utilisation rates among individuals with panic disorder appear disproportionately high (Barsky et al., 1999; Katerndahl and Realini, 1997, Zane et al., 2003). Panic disorder patients consulted their general practitioner significantly more often than the control group (Simpson et al. (1994). More frequent physician visits, emergency room visits and mental health visits were reported among patients with PD as compared to general medical outpatients (Barsky et al., 1999). Patients seek immediate medical care during initial panic attacks, commonly in the Emergency Department (ED), because these attacks are often interpreted as evidence of a catastrophic physical disorder, such as a myocardial infarction (Tueth, 1997; Dammen et al., 2001) or respiratory failure and uncomfortable feelings related to fear of death. This is the reason why patients who screen positive for panic disorder use emergency medical services and ED services more frequently (Zane et al., 2003) than the general population. Although only 60% of people with panic disorder seek care, 32% of these patients present to EDs (Zun, 1997, Fleet et al., 1996). Approximately 35% of patients with PD visit their general family practitioner and continue to receive treatment with their primary physician, while only 26% of panic attack patients initially seek care in a mental health setting (Katerndahl and Realini 1995).

2.2. Risk factors of panic disorder

The pathogenesis of PD is complex and involves sociodemographic, psychological, biological and evolutionary factors. Marital status is a significant risk factor for PD: the highest lifetime prevalence rates are found in widowed, separated or divorced subjects (Wittchen and Essau 1993). In particular, about 25% of subjects with PD had a previous divorce, compared to approximately 10% of those with no active phase of disorder (Yates, 2009). There are no consistent findings concerning the educational level and the risk of developing PD, although Eaton et al. (1994) reported a tenfold higher risk for persons with less than 12 years of education. A high frequency of stressful life events also often precedes the onset of PD (Scocco et al., 2007). Several studies have suggested that life events such as early parental loss or childhood abuse may increase the risk of PD (Faravelli 1995; Tweed et al., 1989; Cogle et al., 2010, Goodwin et al., 2005b, Roy-Byrne et al., 1986; Assellmann et al., 2017). Furthermore, children with behavioural inhibitions may appear to have an increased risk for later development of PD (Reznick et al., 1992; Hirshfeld-Becker et al., 2008).

According to the cognitive theory of Clark (1986), individuals who experience recurrent panic attacks have a relatively enduring tendency to interpret certain bodily sensations in a catastrophic fashion. Barlow (1988) described panic as the basic emotion of fear, which is considered to be an acute reaction to perceived imminent danger when no danger is present. The false suffocation alarm theory by Klein (1993) suggested that PD patients have a low stimulation threshold of the asphisiostat, a physiological mechanism of protection from potentially lethal stimuli. One important vulnerability factor for anxiety disorders, and in particular for PD, is anxiety sensitivity or fear of anxiety symptoms. Studies support the idea that anxiety sensitivity acts as cognitive predisposition for the development of PD (Schmidt et al., 1997). Gorman et al. (1989) were the first authors to propose a neuroanatomical model specific to PD. This aetiological model of PD suggested an abnormal sensitivity in the brain mechanism of a fear and alarm response involving a network of neuronal pathways and multiple neurotransmitter systems. According to this hypothesis, panic attacks originate from a dysfunction in the brain fear network that integrates various structures of the brainstem, hippocampus, amygdala, medial hypothalamus, thalamus and cortical regions. This model links acute panic attacks to the brain stem nuclei, anticipatory anxiety to limbic activation and kindling, and phobic avoidance to prefrontal cortical function, integrating biological and psychological aspects of PD (Gorman et al., 2000).

2.3. Pharmacotherapy of panic disorder and treatment outcome

Five classes of medication have been shown in randomised trials to be more effective than placebo in patients with panic disorder: selective serotonin-reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), high-potency benzodiazepines, tricyclic antidepressants and monoamine oxidase inhibitors (Mitte, 2005; Bradwejn, 2005; Otto et al., 2001; Gould et al., 1995; Susman and Klee 2005; Goddard et al., 2001). A meta-analysis compared the efficacy of three pharmacologic classes of medications with placebo in patients with panic disorder as measured by levels of global anxiety (frequency of panic attacks, and agoraphobia) and depression: SSRIs (in 17 trials), tricyclic antidepressants (in 23 trials) and benzodiazepines (in 25 trials) (Mitte, 2005). The three classes of medication were equally effective in treating anxiety, but the effect of benzodiazepines in treating depression was marginally less than that of either tricyclic antidepressants or SSRIs (Mitte, 2005). Large studies have shown a clinically significant response (defined by a 50% decrease in the frequency of panic attacks or global anxiety) in 50 to 80% of patients treated with SSRIs, tricyclic antidepressants or benzodiazepines (Mitte, 2005). An earlier meta-analysis compared the effects of SSRIs or placebo in 12 randomised, controlled trials (Otto et al., 2001). SSRIs were significantly more effective than placebo in reducing global anxiety and the frequency of panic attacks; more than 50% of patients treated with SSRIs became panic-free in seven of nine studies reporting this outcome. In a large, placebo-controlled trial in patients with panic disorder, SNRI venlafaxine, at a dose of 75 to 225 mg per day, reduced the global severity of panic, anticipatory anxiety, and fear and avoidance of social activities on the basis of validated anxiety scales. However, the drug did not increase the likelihood of becoming free of panic (Bradwejn et al., 2005). Because of their safety profile, as compared with the safety profiles of tricyclic agents and monoamine oxidase inhibitors, SSRIs are recommended as the first drug option in the treatment of panic disorder (Otto et al., 2001, Working Group on Panic Disorder 1998). Because clinical experience suggests that many patients with panic disorder are hypervigilant regarding side effects, SSRIs should be started at low doses, with dose titration every five to seven days, as tolerated. Although the benzodiazepines continue to have an important role in the treatment of panic disorder, concern with respect to dependence, medication abuse, side effects, and the rapid re-emergence of symptoms after discontinuation have led to the recommendation that these agents should not be the first choice for treatment, but instead used “as needed” (Susman and Klee 2005).

The goal of treatment should be to eliminate panic attacks, if possible, because a partial response often results in continued avoidance of frightening situations and impairment in social functioning. Unfortunately, not all panic patients achieve remission, so under-treatment of PD remains a major problem (Lydiard,

2011). The literature review shows that 20–40% of PD patients treated pharmacologically are unresponsive to therapies, similar to the 30–40% of patients treated with cognitive-behavioural therapy (van Apeldoorn et al., 2008). Even the integrated therapies are not able to substantially reduce the percentage of insufficient response in PD (van Apeldoorn et al., 2008). In short-term clinical trials, despite the different treatment options, 17–64% of participants with PD did not respond adequately to pharmacotherapy and continued to have PAs and/or exhibited negative behaviour related to PAs (Freire et al., 2016). Furthermore, follow-up studies demonstrate that 25–50% of patients relapse within 6 months after discontinuation of drug treatment, whereas 40–60% of PD patients show subclinical symptoms after 4–6 years of follow-up (Perna et al., 2012; van Apeldoorn, 2008). At 12-year follow-up, patients with panic disorder without agoraphobia were most likely to show a recovery in all assessment points; however, rates of recurrence were similar for panic disorder both with and without agoraphobia (Bruce et al., 2005). Severe agoraphobic avoidance and the severity of depressive symptoms, particularly the recurrent form, are among the predictors of non-remission and poorer outcomes in PD patients (Bruce et al., 2005; Park et al., 2012; Batelaan et al., 2010, Slaap and den Boer, 2001), along with co-morbidity with generalised anxiety disorder, or social phobia and severity of panic symptoms (Chavira et al., 2009), as well as the coexistence of PD and bipolar disorder (Lee, 2008). On the other hand, PD patients without agoraphobia had higher recovery rates, spent considerably less time in illness episodes and were less affected by comorbid conditions compared with patients with any of the other anxiety disorders (Bruce, 2005).

2.4. Personality and panic disorder

Personality is defined as the unique psychological qualities of an individual that influence a variety of characteristic behaviour patterns (both overt and covert) across different situations and over time (Gerrig and Zimbardo, 2002). Personality traits are defined as highly stable behaviour patterns (Kipper et al., 2009), traditionally conceptualised as having two components: temperament, which refers to biologically based, stable individual differences in emotion and its regulation, and character, which refers to individual differences due to socialisation. However, as the distinctions between these constructs are questionable, the terms “personality” and “temperament” are now often used interchangeably (Caspi and Shiner 2006, Clark and Watson 1999). In recent years, some authors have questioned this stability, suggesting that personality characteristics can be influenced by acute psychiatric symptoms and by time frame (Seviewright et al., 2002, Cohrs et al., 2008). Personality characteristics have been described as dynamic constructs that develop over the lifespan and change in response to maturation and life circumstances (Fraley and Roberts 2005, Rothbart and Bates 2006). A number of processes and factors contribute to stability and a change of personality. For example, genes are considered as a main influence on

phenotypic stability of personality (Krueger and Johnson 2008, Kandler et al., 2010). On the other hand, many of the mechanisms hypothesised to cause personality change (e.g., life events, the timing of various social roles, physical health and cultural values) can differ considerably across cultures (Chopik WJ, Kitayama S. 2018). A recent review showed that personality traits, especially Neuroticism and Extraversion, could change as a result of clinical interventions (Roberts et al., 2017). In contrast to personality traits, which are described in dimensional terms (i.e. less vs. more of a trait), specific personality disorders are instead categorical terms (i.e. presence vs. absence of “disorder”) (Brandes et al., 2006) and represent a severe disturbance in the characterological constitution and behavioural tendencies of the individual, usually involving several areas of the personality, and nearly always associated with considerable personal and social disruption. Personality disorder tends to appear in late childhood or adolescence and continues to manifest into adulthood.

The several models aim to explain the relationship between personality traits and major psychiatric disorders, such as depression and PD (Akiskal et al. 1983, Klein et al., 2011, Clark et al., 1994, Freire et al., 2007, Krueger and Tackett, 2003, Wachleski et al., 2008). In particular, the following models were proposed by Klein et al. (2011):

(a) The continuous/spectrum model, which considers that personality traits and major psychiatric disorders are epiphenomena of the same process, and the relationship between them is not hierarchical. This model proposes that associations between personality and psychopathology are found because these two constructs both occupy a single domain or spectrum, and psychopathology is simply a display of the extremes of normal personality function. Support for this model is provided by an issue of criterion overlap. For instance, two of the primary facet scales of neuroticism are “depression” and “anxiety”. Thus, the fact that diagnostic criteria for depression, anxiety and neuroticism assess the same content increases the correlations between these domains.

(b) The precursor model, in which mental disorders and personality have similar etiology, personality predicts onset of disorder.

(c) The common cause model proposes that Neuroticism and common mental disorders, such as depression and anxiety disorders, are related, though not directly, and they share etiology accounts, genetic and environmental determinants.

(d) The “scar” hypothesis, in which the occurrence of mental disorders, such as PD and depression, could provoke permanent personality alterations, such as increased dependency or insecurity (Akiskal et al., 1983, Ormel et al., 2013). According to the scar model, episodes of a mental disorder ‘scar’ an individual’s personality, changing it in significant ways from premorbid functioning. An example of a scar effect would be a decrease in openness to experience following an episode of post-traumatic stress disorder.

(e) The concomitant model proposes mental disorders shape personality and neuroticism, but the effect of psychopathology is temporary. After the episode has remitted, the effect disappeared.

(f) The pathoplasticity model views personality having a causal influence on depression after onset.

(g) The predisposition or vulnerability model in which personality could facilitate the emergence of mental disorders and is, in fact, part of its etio-pathogenesis. The vulnerability may be general, predisposing the person to a range of distress disorders, or be specific to a particular disorder (Blatt, 1974). High Neuroticism either causes the development of common mental disorders directly or enhances the impact of causal risk factors such as stressful life events.

It is proposed that personality traits may play a crucial role in the onset and development of PD (Clark, Watson, and Mineka, 1994), whereby extremes of personality traits indicate greater dysfunction in patients with anxiety disorders (Brandes et al., 2006). An anxious temperament and anxiety-related personality traits may represent intermediate phenotypes that predispose to PD (Na et al., 2011, de Ruiter, 1992). A strong relationship has been observed between high neuroticism and low positive emotions and PD (Bienvenu et al., 2001b; Bienvenu et al., 2004, Carrera et al., 2006, Zugliani et al., 2017). On the other hand, Introversion was found to be related to agoraphobia, but not to PD itself, while patients with agoraphobia tend to have higher Neuroticism, lower Extraversion or both (Bienvenu et al., 2001a; Bienvenu et al., 2004, Bienvenu et al., 2007). Other personality factors such as Openness, Agreeableness and Conscientiousness seem to be unrelated to PD (Bienvenu et al., 2001b, Enns and Cox, 1997, Carrera et al., 2006). The other reports indicate less Self-Directedness and Cooperativeness (Izci et al., 2014) and more obvious Harm Avoidance and fatigability (Izci et al., 2014; Kennedy et al., 2001; Wiborg et al., 2005) in patients with PD compared to healthy controls. In addition, PD patients show characteristics that are related to a developmental deficit in the capacity to regulate negative affect (Diamond, 1987). In particular, they tended to be shy and inhibited in childhood, especially showing a clear difficulty in expressing aggressiveness (Almeida and Nardi, 2002). The personality traits in PD could be influenced by gender and psychiatric comorbidity. Thus, it was shown that females with PD had higher Extraversion scores and were more fearful of physical consequences of anxiety than males, while males scored higher on angry hostility and depression facets of Neuroticism (Foot et al., 2004). Furthermore, Neuroticism as well as Agreeableness and Extraversion, though not Conscientiousness or Openness, significantly differed in PD patients with one disorder from those with two or more disorders (Cuijpers et al., 2005). Several other studies also showed an accentuation of certain personality traits, including Harm Avoidance, Neuroticism and Extraversion, in PD patients with affective comorbidity as compared to those with PD only (Ampollini et al., 1997; Bienvenu et al., 2001a; Bienvenu et al., 2001b; Freire et al., 2007, Zugliani et al., 2017). Neuroticism and Extraversion have also shown a strong correlation with the accumulation and severity of depression and agoraphobia with PD (Zugliani et al., 2017). Taken together, studies indicate significant alterations in the personality profiles of patients with PD, which are more pronounced when

comorbid with affective disorders. In terms of coexisting personality disorders, the high rates of cluster B (antisocial, borderline, histrionic, and narcissistic) and cluster C (avoidant, dependent, and obsessive-compulsive) personality disorders are reported in PD patients (Yates et al., 2009, Ozkan M, 2005; Skodol et al., 1995; Bienvenu et al., 2009; Bienvenu, 2005, Grant, 2005). The stronger association was observed for borderline personality disorder (BPD, whereas 80% of in-patients with BPD met the criteria of anxiety disorders, including 48% having PD (Zanarini et al., 1998). Notably, that patients having PD with agoraphobia tend to have a higher prevalence of personality disorder diagnoses than those without agoraphobia (Reich, 1987; Iketani et al., 2002).

2.5. Personality and treatment outcome in panic disorder

Although personality traits and disorders are defined as highly stable behaviour patterns (Kipperet al., 2009), several authors have suggested that personality traits can be influenced by acute symptoms and by aging (Rocca et al., 2006, Corchs et al., 2008) or even change, in particular Neuroticism, under the effect of clinical intervention, pharmacotherapy or cognitive-behavioural therapy (Tang et al., 2009, Clark et al., 2003, Barlow et al., 2013).

It should be noted that in contrast to depression the association between personality characteristics and treatment outcome in anxiety disorders, including PD, is less investigated and understood (Bienvenu and Brandes, 2005). However, such a relationship seems to be bilateral where personality may modify response to treatment, but in turn the treatment can lead to certain changes in personality patterns. In particular, it has been shown that personality deviations may affect severity of the PD (Sokol et al., 1995, Ozkan et al., 2005), negatively influence treatment response (Slaap and den Boer 2001; Marchesi et al., 2006a) and contribute to a lesser decrease in specific panic and agoraphobic symptoms during a treatment course in comparison with PD without co-morbid personality disorders (Prasko et al., 2005). However, the evidence that personality disorders may worsen the prognosis of PD or lead to a longer treatment course (Prasko et al., 2005) is not supported by other observations (Hofmann et al., 1998). Controversial results have also been reported regarding the possible effect of personality traits on treatment outcome in PD. Therefore, Navarro et al. (2013) showed that borderline personality traits presenting in PD patients at the onset of the disease might worsen the response to pharmacological treatment. In some other studies, only the borderline traits have negatively influenced the remission of panic attacks following one-year treatment with SSRIs in PD patients, whereas the number of other personality traits did not affect the outcome of medication (Marchesi et al., 2006a; Hoffmann et al., 1998). In contrast, the borderline traits predicted a better treatment outcome at the six-month follow-up observation, whereas obsessive-compulsive personality traits were negatively related to treatment outcome (Dreessen et al., 1994). As was suggested, the effect of personality on treatment outcome in PD may depend on

several factors, including methodological variations across studies, differences in duration of follow-up, applied medications and definition of PD remission (Marchesi et al., 2006a). Earlier, Dreessen et al. (1998) reviewed 15 studies in aiming to evaluate the influence of personality disorders on treatment outcome in anxiety disorders, including PD. They concluded that the short-term treatment outcome (up to 3 months) of PD is not convincingly affected by the presence of categorical or dimensional personality disorder variables. Chavira et al. (2009) demonstrated that higher Neuroticism and higher anxiety sensitivity scores at baseline were associated with less probability of clinical improvement in 3 months in primary care patients with PD, which is in line with an earlier observation of Costa and McCrae (1985) from a 3-month follow-up study suggesting that lower Neuroticism scores at baseline predict better clinical improvement. Both prominent Neuroticism and higher harm avoidance seem to be the most important predictors of non-response to therapy with SSRIs, whereas higher Self-Directedness, Agreeableness and Cooperativeness predict a better response to drug treatment (Marchesi et al., 2006b). However, associations between basic personality dimensions and PD treatment response are not univocal. None of the explored personality traits as assessed by the Neuroticism-Extraversion-Openness Five-factor Personality Inventory was related either to clinical severity or to short-term response to SSRI treatment (Carrera et al., 2006), and no association between personality traits and PD symptom remission were found in other studies (Marchesi et al., 2006a).

In terms of the treatment effect on personality patterns in PD, some first reports have indicated that a large improvement in panic symptoms after anxiolytic medication with alprazolam or diazepam is accompanied by significant changes in neuroticism-related personality traits (Reich et al., 1991; Noyes et al., 1986). A positive influence on personality disorder characteristics in PD was also observed during both individual CBT and imipramine treatment; however, the efficacy was predicted by temperament (Hofmann et al., 1998). A more recent relatively small prospective study found that improvement in the symptoms of PD after 1 year treatment with SSRIs was associated with the normalisation of paranoid traits and persistence of avoidant and dependent characteristics, suggesting that some personality traits in PD show, at least in part, a state phenomenon (Marchesi et al., 2005). In the later study by Marchesi et al. (2008,) higher levels of harm avoidance were found after 1 year of pharmacological therapy in patients with PD in remission compared to healthy controls. A recent study that used the SSP scale to evaluate the effect of 12-weeks of internet-based cognitive behaviour therapy on severe health anxiety has shown a significant reduction on neuroticism related scales, whereas traits relating to extraversion and aggression remained largely unchanged (Hedman et al., 2014). On the contrary, a significant decrease was observed in all anxiety-related scores as well as in the aggression and hostility-related scores on the Karolinska Scales of Personality (KSP) in PD patients after 6 months of treatment with SSRI citalopram (Neuger et al., 2002). Additionally, a significant decrease in the score on 8 of the 10 Minnesota Multiphasic Personality Inven-

tory scales was observed in PD patients following pharmacological treatment. However, the anxious and neurotic personality characteristics were still more pronounced in asymptomatic patients as compared to healthy controls (Kipper et al., 2009). Notably, although Brandes et al. (2006) revealed that remission from PD is generally associated with partial “normalization” of personality traits, some others assumed that personality traits remain stable over the course of PD (Morey et al., 2010, Santor et al., 1997).

3. RATIONALE FOR THE STUDIES

The relationship between personality and PD is still poorly understood and has become a topic of scientific debate and scrutiny. Considering that most of the studies cited above included relatively small samples and provided only preliminary or controversial evidences, further extensive investigations are warranted to evaluate the involvement of personality disposition in the course and treatment of PD. Furthermore, there is a vast number of studies that have sought to identify predictors of treatment response in anxiety disorders, including PD. Although, these studies have not as yet yielded findings that were robust enough to be clinically relevant, there is a strong belief that predictors of treatment response would contribute to the personalised medicine approach and, particularly, guide decision making in selection of the most suitable medication for individual patients. Moreover, similarly to other putative predictors, including demographic and clinical characteristics, personality traits are easy to evaluate and not as costly as genetic or neuroimaging biomarkers, which significantly facilitate the implementation of personality assessment in routine practice. In particular, a better understanding of the relationship between personality traits, a course of PD and its treatment outcome may significantly improve the prediction and management of this prevalent and highly disabling anxiety disorder. Furthermore, personality traits seem to play a crucial role in the pathogenesis and prognosis of all mental health disorders. Therefore, the further exploration of personality traits in patients with mental disorders, including PD, would have an important impact on fundamental knowledge about the development and course of these diseases. In particular, to achieve the goals of the current work, we chose the Swedish universities Scales of Personality (SSP) to measure the personality traits in PD patients. The SSP is a self-rated questionnaire based on the Karolinska Scales of Personality (KSP), an instrument designed to measure stable personality traits related to psychopathology (Schalling 1978; Gustavsson et al., 2000). The KSP scales tap personality traits that have biological correlates and could contribute to the development of mental disorders (Schalling et al., 1987; Klinteberg et al., 1987). Although the KSP does not aim to cover all aspects of personality, it characterises four fairly general temperament dimensions: anxiety, extraversion, socialisation and aggression (Schalling et al., 1987). During development of the SSP, some aggression and neuroticism-related scales of the KSP were dropped due to unsatisfactory psychometric properties. The final version of the SSP comprised 13 scales that demonstrated good psychometric properties in the Swedish normative sample (Gustavsson et al., 2000). Four scales of the SSP assess various aspects of vulnerability to anxiety. Another five scales reflect aggression and related traits. The remaining scales characterise sensation seeking, impulse control, relation to the social environment and conformity. Factor analysis suggests that the SSP scales measure three broader constructs: neuroticism, extraversion and aggressiveness (Gustavsson et al., 2000). The

SSP has shown to be useful as a personality measure in psychobiological research (Damberg et al., 2003, Jönsson et al., 2003). A widely accepted approach in the field of personality structure research is the five-factor model (Costa and McCrae 1992b), which proposes that most individual differences in personality can be understood in terms of five basic dimensions. Relating the personality traits measured by the SSP with the five-factor model could give a better understanding of the position of these traits in the basic structure of personality. One of the standard instruments designed to measure these five factors and their facets is the Revised NEO Personality Inventory (NEO-PI-R). The NEO-PI-R has demonstrated a similar structure in various cultures (McCrae and Costa 1997). The SSP intends to measure temperament-like features, meaning that it also should be applicable in different social and cultural contexts. At the same time, the reliability and factor structure of the SSP in samples with different language and cultural backgrounds have not yet been studied. Thus, we aimed to test the reliability and validity of the Estonian version of the SSP. This also allows us to investigate the universality of the personality factors assessed by the SSP in a different social and cultural background. We also intend to characterise the position of SSP-measured traits within the basic personality dimensions of the five-factor model. Taken together, we found the SSP to be a promising instrument for its clinical exploitation in the evaluation of personality traits among patients with PD.

4. AIMS OF THE STUDIES

The general objective of the present work was to characterise the personality traits of patients with PD and to explore the relationship between personality disposition and treatment outcome in those patients by using validated SSP assessment. The specific aims were, as follows:

1. To test the reliability and validity of the Estonian version of the SSP (I)
2. To characterise the position of SSP-measured traits within the basic personality dimensions of the five-factor model (I)
3. To identify the differences in personality traits evaluated by SSP between patients with PD and healthy subjects (II)
4. To examine differences in personality dimensions between PD patients with and without affective comorbidity, and explore the relationship between SSP personality domains and various demographic and clinical variables of PD (II)
5. To examine the effect of escitalopram treatment on personality traits in PD patients (III)
6. To identify whether the treatment outcome of PD could be predicted by any personality trait as measured by using SSP (III)

5. MATERIALS AND METHODS

5.1. Subjects

A total of 529 subjects participated in the study I, among them 174 (32.9%) men and 355 (67.1%) women. The average age was 35.7 years (standard deviation, $s=14.0$ years, range 18–74). The study group was formed of two samples. Sample 1 consisted of 331 healthy volunteers (106 men, 225 women, average age 39.1, $s=13.5$) who participated in different research projects conducted at the Department of Psychiatry of the University of Tartu. The absence of psychiatric disorders in this sample was confirmed by the Mini International Neuropsychiatric Interview (M.I.N.I. 5.0.0; Sheehan et al., 1998). Sample 2 was a convenience sample comprising 198 subjects with various educational and social backgrounds (68 men, 130 women, average age 30.1 years, $s=12.9$). The sample consisted of university students, their acquaintances, relatives and workmates.

The study II sample consisted of 193 patients (45 men, 148 women, average age 36.9, $s=12.3$) with PD recruited at the Psychiatry Clinic of the Tartu University Hospital and 314 healthy subjects (94 men, 220 women, average age 38.9, $s=13.8$) recruited by newspaper advertisement in Tartu, Estonia. Diagnosis of PD according to DSM-IV criteria was verified using the Mini International Neuropsychiatric Interview (M.I.N.I. 5.0.0; Sheehan et al., 1998) and substantiated by psychiatric history and medical records. At the time of assessment, 126 of patients (65.3%) had current symptoms of PD and 67 were in remission. PD patients with current or past comorbidity with mood disorders or with other anxiety disorders were included in the study, but no other psychiatric comorbidity was allowed. Personality disorder was not assessed, it may be considered as limitation of the study. According to the clinical assessments 71 (36.8%) patients were defined as “PD only” group, who never met criteria for any other psychiatric diagnosis, with the exception of agoraphobia. The remaining 122 patients (63.2%) with comorbid mood or anxiety disorders were defined as “PD comorbid” group. Major depressive disorder was the most frequent comorbid or lifetime co-existing condition, which was diagnosed in 93 patients (with its current presence in 60 patients); whereas the criteria for bipolar disorder were met in 29 patients (with its current presence in 7). Other anxiety disorders, including social phobia and obsessive-compulsive disorder, were detected in 46 patients. Concurrent agoraphobia was present in 115 (59.6%) of the patients, among them 46 were from “PD only” and 69 from “PD comorbid” group. The earlier onset of PD with first panic attacks before age of 30 years was observed in 106 patients, whereas in 87 patients PD has developed in later lifetime period. The healthy subjects were interviewed using the M.I.N.I. and questioned about their family psychiatric history. Only those without personal or family (defined as first-degree relatives) history of psychiatric disorders were included in this study. The majority of the subjects were of Estonian ethnicity with a

similar between-group distribution (99% among patients and 95% among controls).

The study III sample consisted of 110 adult patients with PD. Participants were recruited at the Psychiatry Clinic Outpatient Department or through referral from the Emergency Department of the North Estonia Regional Hospital. Three patients were excluded (two by patient request, one lost to follow-up); therefore, 107 PD patients (mean age = 34, s = 11.7 years; 71 females) were included in the final sample of the study. The diagnosis of PD according to the DSM-IV criteria was verified using the Mini International Neuropsychiatric Interview and substantiated by psychiatric history and medical records. PD patients with current or past co-morbidity with mood disorders or with other anxiety disorders were allowed to take part in the study. However, other problems, such as schizophrenia or another psychotic disorder, severe suicide risk, substance abuse or dependence, organic mental disorder, severe personality disorder, serious unstable medical condition (i.e. endocrinological, liver, respiratory, cardiovascular diseases), were excluded. The majority of the subjects were of Estonian ethnicity (84.1%). All patients were unmedicated and did not receive any formal psychological help for at least 3 months before this trial and during the investigation.

5.2. Ethical considerations

The Human Studies Ethics Committee of the University of Tartu approved the study protocols, and all participants provided written informed consent.

5.3. Measures

5.3.1. The Mini International Neuropsychiatric Interview (M.I.N.I.5.0.0.)

Diagnosis of PD and psychiatric co-morbidity according to DSM-IV criteria was verified using the Mini International Neuropsychiatric Interview (M.I.N.I.5.0.0.; Sheehan et al., 1998) and substantiated by psychiatric history and medical records (II, III). The Mini International Neuropsychiatric Interview (M.I.N.I.) is a structured diagnostic interview, developed for DSM-IV and ICD-10 psychiatric disorders. In 1999 M.I.N.I. 5.0.0. was adapted to Estonian at the Department of Psychiatry of Tartu, University of Tartu (Shlik et al., 1999). The absence of psychiatric disorders in sample of healthy subjects in control group (I, II) was confirmed by the M.I.N.I.5.0.0.; also subjects were questioned about their family psychiatric history (II).

5.3.2. Swedish universities Scales of Personality (SSP)

The Swedish universities Scales of Personality (SSP) is a self-rated questionnaire based on the Karolinska Scales of Personality (KSP) designed to measure stable personality traits related to psychopathology (Gustavsson et al., 2000). The SSP comprises of 91 items grouped into 13 scales: Somatic Trait Anxiety (STA), Psychic Trait Anxiety (PsTA), Stress Susceptibility (SS), Lack of Assertiveness (LA), Impulsiveness (I), Adventure Seeking (AS), Detachment (D), Social Desirability (SD), Embitterment (Em), Trait Irritability (TI), Mistrust (M), Verbal Trait Aggression (VTA), and Physical Trait Aggression (PHTA). Each scale is formed by 7 items rated on a scale of 1 = does not apply at all, to 4 = applies completely.

The factor analysis of the original SSP yielded three factors: neuroticism, aggressiveness and extraversion (Gustavsson et al., 2000).

The raw scores were converted into T-scores using the raw score means and standard deviations of men and women from Estonian normative sample in study I. Only T-scores were used in the analysis in studies II and III.

All participants self-evaluated their personality traits using the Estonian version of SSP (I, II, III) (Aluoja et al., 2009).

- (I) The SSP was completed by all participants of the study (both samples).
- (II) At the time of assessment all the patients (with current symptoms of PD and in remission) and all the control subjects completed the Estonian version of SSP to evaluate their personality traits.
- (III) The Estonian version of SSP was completed by study group twice, at the beginning of the study and at the last visit after 12-week treatment with SSRI escitalopram.

5.3.3. The revised NEO Personality Inventory (NEO-PI-R).

The NEO-PI-R is a 240-item self-report questionnaire that measures personality structure according to the five-factor model (Costa&McRae, 1992b). Similarly to the original, the Estonian version of the NEO-PI-R (Kallasmaa et al., 2000) provides scores on the five personality dimensions: Neuroticism (N), Extraversion (E), Openness (O), Agreeableness (A), and Conscientiousness (C). Each personality domain is composed of six facet scales. Responses are made on a five-point Likert-type scale (from strongly agree to strongly disagree). One of the standard instruments designed to measure these five factors and their facets is the Revised NEO Personality Inventory (NEO-PI-R) (Costa et al., 1992). NEO-PI-R has demonstrated similar structure in various cultures (McCrae&Costa 1997). In study I 197 subjects from the Sample 2 completed Estonian version of the NEO-PI-R.

5.3.4. Panic Disorder Severity Scale (PDSS)

The PDSS (Shear et al., 1997) is used for assessment of clinical severity and treatment response of PD. The PDSS is seven-item measure, score range 0–28, comprised of five items assessing core DSM-IV symptoms of PD, with or without agoraphobia, and two items rating work and social impairment (Shear et al., 1997). The scale has been considered sensitive to change with medication treatment (Pollack et al., 1998). The scale has been translated into several languages and findings strongly support the reliability and validity of the PDSS in different cultural background (Shear et al., 2001, Lee et al., 2009, Barlow et al., 2000). For screening of patients a cut-off score 8 was recommended, which identifies patients with current PD with a sensitivity of 83.3% and a specificity of 64% (Shear et al., 2001). The PDSS score 7 or less has been used in several studies for defining of remission (Furukawa et al., 2009, Shear et al., 2001, Park et al., 2012, Seki et al., 2016) and is considered the point of remission in this study. All patients in the study III were assessed using the PDSS beginning from the baseline up to the endpoint of the study in every two weeks.

5.3.5. The Clinical Global Impression scale (CGI)

The Clinical Global Impression scale (Guy, 1976) is a 3-item observer-rated scale commonly used to measure symptom severity, global improvement, and therapeutic response. Each component of the CGI is rated separately. The Clinical Global Impression – Severity scale (CGI-S) is a 7-point scale to rate the severity of patient illness at the time of assessment, relative to the clinician's past experience with patients who have the same diagnosis. Considering total clinical experience, a patient is assessed on severity of mental illness at the time of rating 1, normal, not at all ill up to 7, extremely ill.

The Clinical Global Impression – Improvement scale (CGI-I) is a 7 point scale for assessment how much the patient's illness has improved or worsened relative to a baseline state at the beginning of the intervention and rated as: 1, very much improved; 2, much improved; 3, minimally improved; 4, no change; 5, minimally worse; 6, much worse; or 7, very much worse.

The Clinical Global Impression – Efficacy Index is a 4 point \times 4 point rating scale that assesses the therapeutic effect of the treatment as 1, unchanged to worse; 2, minimal; 3, moderate; 4, marked by side effects rated as none, do not significantly interfere with patient's functioning, significantly interferes with patient's functioning and outweighs therapeutic effect. With this scale therapeutic efficacy and treatment-related adverse events should be taken into account. Clinical severity and treatment response in the study III were assessed using the CGI, every two weeks during the study.

5.3.6. Hamilton Anxiety Scale (HAM-A)

Hamilton Anxiety Scale (HAM-A) (Hamilton, 1959) was developed to measure the severity of anxiety symptoms and is widely used in clinical and research settings. The HAM-A is an interview-based scale, consists of 14 items, each defined by a series of symptoms, and measures psychic anxiety (mental agitation and psychological distress) and somatic anxiety (physical complaints related to anxiety). Each item is simply given a 5-point score – 0 (not present) to 4 (severe). HAM-A was used as a secondary scale in the study III for clinical assessments, every two weeks during the study.

5.3.7. Toronto Side-effect Scale (TSES)

Toronto Side-effect Scale (TSET) is a 32-item instrument used to evaluate possible adverse events, incidence, frequency, and severity of central nervous system, gastrointestinal, and sexual side effects (Vanderkooy et al., 2002). TSES was used as a secondary scale for clinical assessments in the study III, every two weeks during the study.

5.3.8. The Emotional State Questionnaire (EST-Q)

The Emotional State Questionnaire (EST-Q) (Aluoja et al., 1999) is a self-rated questionnaire, constructed on the basis of the relevant diagnostic criteria in the DSM-IV (APA, 1994) and the 10th revision of the International Classification of Diseases and Related Health Problems (ICD-10, World Health Organization 1993). The EST-Q contains 28 questions in five subscales; items are rated on a 5-point frequency scale ranging from 0 (never) to 4 (continuously).

5.3.9. Sociodemographic measures

Data on sex, gender, age, years of education, employment, smoking habit, alcohol drinking frequency, presence of somatic co-morbidities and marital status were recorded in the study groups according to the study design (I, II, III) from the structured interviews.

5.4. Treatment

The patients in the study III were treated with 10–20 mg/day of escitalopram for 12 weeks using an open-label placebo non-controlled study design. No other medications were permitted during the treatment period, except for hormonal contraceptives. Additionally, zolpidem or zopiclon for insomnia and alprazolam

for acute anxiety or panic symptoms were also allowed during the first 6 weeks. All patients started the treatment with a dose of 10 mg/day of escitalopram for the first 4 weeks. The patients who showed at least a 50% decline in the PDSS total score at week 4 continued taking 10 mg of escitalopram until the end of the study. The dose of escitalopram was increased and maintained at 20 mg in patients who demonstrated less than a 50% decrease in PDSS total score at week 4. At the end of week 12, remission was defined as a patient who meets all three criteria – the score on the CGI improvement scale 2 or less, the PDSS score 7 or less – the cut-off point, used in several studies for defining of remission, and no panic attacks for at least the last 2 weeks. Patients who did not meet these criteria were defined as non-remitters. The severity of the panic symptoms and treatment response were rated by a psychiatrist (UV), who was blind to the personality trait measures. During each visit, the patients were also asked to report on their regularity of taking the medication. Adherence to medication was good – by patients self-report none of the patients had more than 3 days drug holiday.

5.5. Statistical analysis

The data were analyzed using the software package STATISTICA 8 (StatSoft Inc., Tulsa, OK, USA) and statistical software R 3.1.1. (R Core Team, 2014). In the study I the internal consistency of the SSP scales was assessed by calculating Cronbach's α coefficients. To evaluate the factorial validity of the SSP a principal component analysis with a varimax rotation was conducted on the scales. The factors were extracted on the basis of an eigenvalue greater than one. In subsample 2 the external validity of the SSP was evaluated by calculating Pearson product moment correlations between the SSP scales and the NEO-PI-R scales. Correlations between the SSP and NEO-PI-R scale facets were reported only if they provided additional information for understanding of the SSP scales. Student's t-test was used to evaluate sex differences. In the study II the non-parametric tests, including Chi-squared, Fisher's exact test and Mann-Whitney U-test, were used as appropriate. Correlations were estimated with Pearson product-moment correlation analysis. Multiple logistic regressions were used to analyze the relative influence of personality variables. Only variables which showed a significant relationship with PD in univariate analyses were entered into multivariate models. Three broader factors, neuroticism, extraversion and aggressiveness, were included in our analysis. The comparisons of SSP subscales and factors were performed for all patients and separately for subgroups of those with and without comorbidity. The comparisons of SSP traits and factors were also done in patient subgroups divided according to pre-defined clinical phenotypes (current PD versus remission status, presence or absence of agoraphobia, early or late onset of PD, presence or absence of family psychiatric history). The results of between-group comparisons of SSP scores were considered nominally significant at $p < 0.05$ level with a conservative

estimation of significance after Bonferroni correction for multiple comparisons on 13 SSP traits at $p < 0.00384$ level. In the study III baseline and final SSP sub-scales were described by mean, minimum and maximum, Wilcoxon signed rank test was used to test the significance of the change between the end and baseline scores of the study. The association between the baseline scores of SSP sub-scales and the changes of PDSS, CGI-S, HAM-A and EST-Q depression was evaluated using partial correlation coefficients with 95% confidence intervals (CI). Similarly, the association between changes in the scores of SSP sub-scales and changes of PDSS, CGI-S, HAM-A and EST-Q depression was evaluated using partial correlation coefficients with 95% CI. The association between the baseline scores of SSP sub-scales and the probability of good treatment response (final PDSS ≤ 7) was analysed by binary logistic regression resulting in odds ratios (OR) with 95% CI. All of the analyses of associations were adjusted for sex, age, years of education, level of drinking frequency, presence of somatic co-morbidities and for the presence of mild depression and/or agoraphobia. The resulting p -values were considered nominally significant at $p < 0.05$ level; the Holm-Bonferroni correction for multiple comparisons was used. Regression assumptions were checked by graphical methods, where applicable.

6. RESULTS AND DISCUSSION

6.1. Study I: Factor structure of the Swedish universities Scales of Personality and position within the five-factor model in an Estonian sample

6.1.1. The gender differences for the SSP

Mean values of the SSP for the whole study sample and separately for males and females are presented in Table 1. Females scored significantly higher on Psychic Anxiety, Stress Susceptibility, Lack of Assertiveness, and Social Desirability. Men scored higher on Verbal and Physical Aggression, Mistrust, and Adventure Seeking. The highest gender differences emerged for Aggression and Anxiety scales. Internal consistency coefficients for the SSP scales ranged from 0.58 (Detachment) to 0.85 (Trait Irritability).

Table 1. Means, standard deviations, Cronbach alphas, and gender differences for the SSP scales

SSP scales	Total			Men		Women		t
	Mean	s	(α)	Mean	s	Mean	s	
Somatic Trait Anxiety	1.84	0.54	0.75	1.82	0.52	1.84	0.55	0.39
Psychic Trait Anxiety	2.17	0.59	0.82	2.03	0.56	2.24	0.60	4.00‡
Stress Susceptibility	2.17	0.50	0.73	2.11	0.50	2.20	0.49	1.97*
Lack of Assertiveness	2.35	0.48	0.64	2.28	0.48	2.39	0.48	2.64†
Impulsiveness	2.42	0.45	0.62	2.42	0.43	2.42	0.46	0.17
Adventure Seeking	2.63	0.56	0.80	2.71	0.55	2.59	0.56	-2.21*
Detachment	2.23	0.44	0.58	2.27	0.44	2.21	0.44	-1.62
Social Desirability	3.05	0.41	0.66	2.95	0.39	3.09	0.41	3.81‡
Embitterment	1.99	0.49	0.73	2.00	0.51	1.99	0.48	-0.30
Trait Irritability	2.30	0.61	0.85	2.33	0.58	2.28	0.63	-0.99
Mistrust	2.11	0.50	0.76	2.21	0.49	2.06	0.49	-3.34†
Verbal Trait Aggression	2.32	0.59	0.78	2.51	0.57	2.23	0.57	-5.45‡
Physical Trait Aggression	2.10	0.62	0.84	2.44	0.60	1.93	0.57	-9.54‡

* $p < 0.05$, † $p < 0.01$, ‡ $p < 0.001$; Means and standard deviations are presented as raw scores.

6.1.2. The principal component analysis of the SSP scales

Present study yielded three factors in analysis of the SSP scales, with an eigenvalue greater than unity, accounting for 62.9% of the total variance (Table 2). The first factor grouped scales that characterise neuroticism, accounting for 27.8% of the variance. Both forms of Trait Anxiety (somatic and psychic), Stress Susceptibility, Lack of Assertiveness, and Embitterment had highest loadings on this factor. The second factor accounted for 20.2% of the variance and included Verbal and Physical Trait Aggression, Trait Irritability, Detachment, and low Social Desirability, thus indicating aggression and nonconformity. Although Detachment and Trait Irritability had highest loadings on the second factor, they also loaded somewhat on the third and the first factors. The third factor comprised Impulsiveness and Adventure Seeking, accounting for 14.8% of the variance. Mistrust loaded nearly equally on two factors, relating to neuroticism and aggression.

Table 2. Results of the principal component analysis with a varimax rotation of the SSP scales

Scales	Factor 1*	Factor 2*	Factor 3*
Somatic Trait Anxiety	0.68	0.26	0.29
Psychic Trait Anxiety	0.89	0.08	0.01
Stress Susceptibility	0.76	0.25	-0.03
Lack of Assertiveness	0.80	-0.14	-0.09
Impulsiveness	0.20	0.22	0.75
Adventure Seeking	-0.03	0.07	0.79
Detachment	0.38	0.50	-0.40
Social Desirability	-0.14	-0.70	0.04
Embitterment	0.65	0.31	0.20
Trait Irritability	0.49	0.58	0.36
Mistrust	0.51	0.50	0.01
Verbal Trait Aggression	0.05	0.72	0.47
Physical Trait Aggression	0.03	0.70	0.29

* Items defining the factors are boldfaced

6.1.3. The correlations between the SSP and NEO-PI-R scales

The correlations between the SSP scales and the five dimensions of personality assessed by the Estonian NEO-PI-R are shown in Table 3. Due to the large number of correlations performed, Bonferroni adjusted alpha of $p < 0.001$ was applied to correlation coefficients. NEO-PI-R domain Neuroticism correlated mainly with SSP neuroticism factor scales. The strongest relationships were found with Psychic Trait Anxiety and Stress Susceptibility. Extraversion related

positively to Adventure Seeking and Impulsiveness, and negatively to Detachment, Psychic Anxiety, Lack of Assertiveness, Stress Susceptibility, and Mistrust. Agreeableness correlated positively with SSP Social Desirability and negatively to Aggression-Irritability scales, Mistrust, and Embitterment. NEO-PI-R dimensions Openness and Conscientiousness had somewhat weaker correlations with the SSP. Conscientiousness related highest to Social Desirability, and Openness to Adventure Seeking. Two SSP scales, Impulsiveness and Physical Trait Aggression had only one moderate correlation with NEO-PI-R domains. To get a better understanding of the relationships of SSP scales with five factors we also looked at the correlations with NEO-PI-R facet scales. SSP Impulsiveness correlated positively with Neuroticism facets Impulsiveness ($r=0.44$) and Hostility ($r=0.32$), Extraversion facets Assertiveness ($r=0.29$), Activity ($r=0.24$), Excitement-Seeking ($r=0.31$), and Positive Emotions ($r=0.29$), and negatively with Agreeableness facets Compliance ($r=0.32$) and Modesty ($r=-0.31$). The highest negative correlation emerged between Impulsiveness and Deliberation ($r=-0.54$), which is a facet of Conscientiousness. SSP Physical Trait Aggression correlated positively with Neuroticism facet Hostility ($r=0.34$), and negatively with Agreeableness facets Trust ($r=-0.25$), Straight-forwardness ($r=-0.34$), and Compliance ($r=-0.54$).

Table 3. Correlations between the SSP and the NEO-PI-R scales in sample 2 ($n=197$)

	N	E	O	A	C
Somatic Trait Anxiety	0.61*	-0.20	-0.12	-0.24	-0.23
Psychic Trait Anxiety	0.76*	-0.48*	-0.25*	-0.04	-0.23
Stress Susceptibility	0.67*	-0.44*	-0.19	-0.10	-0.38*
Lack of Assertiveness	0.44*	-0.45*	-0.27*	0.19	-0.17
Impulsiveness	0.13	0.35*	0.10	-0.19	-0.19
Adventure Seeking	-0.03	0.53*	0.32*	-0.16	-0.02
Detachment	0.30*	-0.67*	-0.35*	-0.20	-0.13
Social Desirability	-0.35*	0.21	0.02	0.52*	0.46*
Embitterment	0.65*	-0.22	-0.23	-0.32*	-0.35*
Trait Irritability	0.55*	-0.12	-0.08	-0.50*	-0.28*
Mistrust	0.48*	-0.36*	-0.26*	-0.43*	-0.18
Verbal Trait Aggression	0.27*	0.10	0.15	-0.60*	-0.27*
Physical Trait Aggression	0.16	0.02	-0.06	-0.44*	-0.07

* $p<0.001$

6.1.4. Discussion

The aim of this study was to test the reliability and validity of the Estonian version of the SSP in order to use the scales in further studies of the role of personality factors in mental disorders. Thus, we evaluated the Estonian SSP in

reference to the Swedish original data, and we characterised the position of the SSP-measured traits within the basic personality dimensions of the five-factor model. The internal consistency of the Estonian SSP scales was in an acceptable range and resembled the Swedish normative data (Gustavsson et al., 2000). The range of the Cronbach α coefficients was between 0.59 and 0.84 in the Swedish study, while they ranged from 0.58 to 0.85 in the Estonian sample. The internal consistency of the anxiety and aggression scales was good and comparable with the original sample. Only two scales, Detachment and Impulsivity, had somewhat low internal consistency. General trends in sex differences were similar in the Estonian and Swedish samples: women scored higher on neuroticism-related scales and men on aggression scales. The one exception was not finding gender differences in the Detachment scale, while in Swedish KSP and SSP studies men have consistently scored higher on detachment than women (Gustavsson et al., 2000; Gustavsson 1997).

The Estonian data revealed a similar three-factor structure of the SSP that was found in the Swedish sample. The Neuroticism factor was identical in the two studies. This finding is important, as clinical researches have identified neuroticism as one of the key contributors to psychopathology (Lahey, 2009). Although the SSP was constructed more for the assessment of psychopathology and the NEO-PI-R for normative personality, both contain Neuroticism factors that are strongly correlated. This also confirms the universality of the general Trait of Neuroticism.

We found analogous factors to Aggression and Extraversion of the Swedish sample (Gustavsson et al., 2000), except for Detachment scale, which according to the Swedish data loaded negatively into the extraversion factor but in our study was closer to aggression. An ambiguous nature of Detachment scale has been earlier observed in factor-analytical studies of the KSP. Some studies indicated that Detachment was a sign of low sociability and thus related to Extraversion (Schalling et al., 1987, Zuckerman et al., 1991), whereas other studies showed its relatedness to Irritability, low Social Desirability and Suspicion (Gustavsson et al., 1997; Ortet et al., 2002). These results probably reflect the complex nature of the construct of Detachment, which is a tendency to avoid close contacts but is also related to the perception of others as being hurtful or untrustworthy (Bornstein et al., 2003). The latter aspect is related to an affective-cognitive side of aggression (irritability and hostility), but not necessarily to its behavioural component represented by Verbal and Physical Aggression scales of the SSP. It is of interest that Detachment and Irritability may even have some common biological correlates (Stalenheim, 2004).

The Neuroticism factor of the SSP is represented in major models of personality (Costa and McCrae, 1992b, Eysenck and Eysenck 1985). The same holds for the second SSP factor that was earlier conceptualised as Extraversion, and combines Adventure Seeking with Impulsiveness (Gustavsson, 2000, Schalling et al., 1987). The third SSP factor, Aggression, joins tendencies to act aggressively with irritability and unfriendly attitudes, and thus resembles Eysenck's Psychoticism (Eysenck and Eysenck 1976). This is in line with other

studies confirming that readiness to express aggression and hostile attitudes form a separate personality dimension (Zuckerman et al., 1991; Zuckerman et al., 1993).

The relationships of the SSP factors with basic personality dimensions were also confirmed by the correlations with the NEO-PI-R scales. Overall, the relationships with the five-factor model were in the expected direction. Scales belonging to the SSP Neuroticism factor had the strongest correlations with the NEO-PI-R Neuroticism scale, confirming that they assess a tendency to experience negative emotions, especially anxiety. Impulsiveness and Adventure Seeking were joined in the same factor and had somewhat similar correlates of NEO-PI dimensions. Both were related to Extraversion but Adventure Seeking was also connected to Openness. This suggests that the need for change and action is related to curiosity and openness to ideas. It has been demonstrated that NEO-PI Openness is related to sensation seeking (García et al., 2005), which is analogous to the SSP Adventure Seeking. The SSP Impulsiveness deserves closer inspection. Impulsivity is a construct that has been positioned differently in the structure of personality. In the five-factor model, it has been placed in the Neuroticism domain (Costa and McCrae, 1992b), while Zuckerman et al. (1991) considered Impulsiveness the closest to sensation seeking and distinct from Neuroticism. Our results show that the SSP Impulsiveness had the strongest association with the low Deliberation of Conscientiousness domain. Similarly, a previous study of health-related personality traits found a relationship between the SSP Impulsiveness items and the Conscientiousness construct (Gustavsson et al., 2003). This indicates that from several aspects of impulsivity, the SSP Impulsiveness measures a lack of premeditation, i.e. a tendency to act on the spur of the moment and without considering the consequences. Like the SSP, most existing impulsivity measures assess lack of premeditation, which has been associated with NEO-PI-R low Deliberation (Whiteside and Lynam, 2001). This aspect of impulsivity is also related to several forms of psychopathology and is closest to Dickman's concept of dysfunctional impulsivity (Dickman, 1990). These results might give a new perspective to the SSP factor combining Impulsiveness with Adventure Seeking. Often in KSP and SSP studies, this factor has been regarded as reflecting the extraversion construct (Gustavsson et al., 2000; Schalling et al., 1987), but present results suggest that the psychological core of it could be a lack of constraint or disinhibition rather than sociability and energy. Clark and Watson (Clark and Watson, 1999) have argued that disinhibition versus constraint is the third general temperament factor besides negative emotionality/neuroticism and positive emotionality/extraversion. Further studies are needed to confirm whether the combination of non-planning impulsivity and adventurousness reflects extraversion or disinhibition, but the latter could be in better agreement with the aim of the SSP in assessing traits constituting vulnerability to psychiatric disorders.

The SSP aggression scales are associated with the NEO-PI-R low Agreeableness and high Neuroticism. In the five-factor model, aggression belongs to

several dimensions, such as Neuroticism that contains the Angry hostility facet, and Agreeableness, concerning the Compliance facet in particular (Costa and McCrae, 1992). Our results are in accordance with Martin et al. (2000) showing that the behavioural, affective and cognitive components of aggression have different NEO-PI-R correlates. Our study supports the idea that the tendency towards aggressiveness constitutes a separate general personality dimension. At least when considering personality traits related to psychopathology, it seems appropriate to treat aggressiveness as a separate construct. Overall, the relationships between the NEO-PI-R and the SSP confirm the validity of the latter instrument. The discrepancies between the two instruments may result from the fact that the SSP has more specific orientation than the NEO-PI-R. While SSP tries to capture nuances of character pathology, such as different types of aggression and anxiety, the NEO-PI-R aims to give a comprehensive assessment of normal personality traits.

A limitation of the study should be addressed. Our study did not include randomly selected population sample like the Swedish study, but instead consisted of healthy research volunteers and a convenience sample. It has been argued that volunteers in research projects cannot be considered a normative sample because of the high rate of psychopathology among them (Shtasel et al., 1991). We tried to correct that possible bias by using a structured diagnostic psychiatric interview (MINI) to rule out mental disorders in the sample of volunteers.

6.2. Study II: Evaluation of personality traits in PD using SSP

6.2.1. Demographic and clinical characteristics

There were no significant differences between patient and control groups in age and sex distribution or in other demographic characteristics, including educational level, marriage and occupational status, however more smoked cigarettes per day were reported by patients as compared to controls (Table 4). None of demographic variables significantly differed between “PD only” and “PD comorbid” groups. Both patient groups had similar mean age of onset of PD ($p=0.49$) and similar proportions of those with or without agoraphobia ($p=0.26$), with current panic symptoms or in remission ($p=0.67$) or of those with or without somatic comorbidity ($p=0.50$). However patients with comorbid PD had a higher load of family history of psychiatric disorders than “PD only” group (51.3% vs. 33.8%, respectively; $p=0.019$), and more smokers were found in comorbid PD group than in pure PD (41.8% vs. 26.8%, $p=0.036$).

Table 4. Demographic characteristics of patient and control groups

Characteristics	PD patients	Healthy subjects	Statistics
Females	148 (76.7%)	220 (70.1%)	
Males	45 (23.3%)	94 (29.9%)	$\chi^2=2.63, p=0.11$
Mean age	36.9 \pm 12.3	38.9 \pm 13.8	F=2.83; $p=0.09$
Education	13.8 \pm 2.9	14.1 \pm 2.9	F=1.29; $p=0.26$
Married or common-law	130 (67.4%)	209 (66.6%)	$\chi^2=0.03, p=0.85$
Employed or student	146 (75.7%)	249 (79.3%)	$\chi^2=0.93, p=0.34$
Smokers	70 (36.3%)	109 (34.7%)	$\chi^2=0.13, p=0.72$
Cigarettes per day	5.4 \pm 8.2	3.7 \pm 6.6	F=5.63; $p=0.02$

6.2.2. Personality differences between total PD group and healthy controls

The analysis of SSP assessments revealed robust differences in personality traits between patients with PD and healthy controls (Table 5). All traits, except for Detachment and Physical Trait Aggression, were significantly deviated in PD group, with the scores for Adventure Seeking and Social Desirability being lower, and all the others being higher in patient group. The SSP factors of Neuroticism and Aggressiveness, but not Extraversion, were significantly higher in PD group than in controls (Table 5). In multivariate logistic regression analyses, Somatic Trait Anxiety, Stress Susceptibility, Lack of Assertiveness and Adventure Seeking remained significant predictors of PD (Table 6).

Table 5. SSP personality characteristics in PD and control groups

SSP traits	PD all (n=193)	PD only (n=71)	PD comorbid (n=122)	Control (n=314)	PD all vs. Control	PD only vs. Control	PD comorbid vs. Control	PD comorbid vs. PD only
Somatic Trait Anxiety	66.8±11.2	63.0±9.8	68.9±11.4	46.6±8.4	<0.000001	<0.000001	<0.000001	0.000249
Psychic Trait Anxiety	60.3±10.8	56.8±11.3	62.4±10.0	47.6±9.6	<0.000001	<0.000001	<0.000001	0.000892
Stress Susceptibility	60.8±10.9	56.7±10.0	63.1±10.7	47.6±9.6	<0.000001	<0.000001	<0.000001	0.000128
Lack of Assertiveness	55.1±11.5	53.2±11.8	56.1±11.2	48.7±9.9	<0.000001	0.004	<0.000001	0.087
Impulsiveness	51.0±10.4	50.3±11.1	51.5±10.1	48.2±9.8	0.003	0.219	0.002	0.309
Adventure Seeking	45.7±10.1	44.9±9.6	46.2±10.3	48.6±10.2	0.002	0.004	0.034	0.352
Detachment	52.0±9.7	49.9±10.3	53.3±9.2	49.5±9.1	0.007	0.810	0.000132	0.005
Social desirability	47.4±9.5	47.9±8.8	47.1±10.0	51.9±9.9	0.000001	0.001	0.000008	0.508
Embitterment	57.0±11.1	52.2±9.5	59.8±11.0	48.8±9.7	<0.000001	0.008	<0.000001	0.000002
Trait Irritability	57.5±9.8	54.4±9.5	59.4±9.5	46.9±9.3	<0.000001	<0.000001	<0.000001	0.0001
Mistrust	53.4±12.0	50.8±10.3	54.8±12.7	48.7±9.5	0.000012	0.068	0.000003	0.067
Verbal Trait Aggression	51.0±9.8	49.4±10.1	51.9±9.6	47.2±9.1	0.000038	0.112	0.000009	0.100
Physical Trait Aggression	50.8±11.0	50.2±11.6	51.2±10.6	47.9±9.5	0.006	0.301	0.003070	0.255
SSP factors								
Neuroticism	60.0±8.9	56.4±8.2	62.1±8.6	47.9±7.3	<0.000001	<0.000001	<0.000001	<0.000001
Extraversion	48.4±8.7	47.6±9.3	48.9±8.4	48.4±8.5	0.935	0.323	0.550	0.188
Aggressiveness	53.0±7.5	51.5±7.2	53.9±7.6	47.5±7.0	<0.000001	0.0001	<0.000001	0.034

Bold *p*-values: the Bonferroni adjusted *p*-value corrected for multiple comparisons on 13 SSP traits at the $p<0.004$ level.

Table 6. Multivariate logistic regression analyses for PD (versus control group)

SSP traits	Total PD group				PD only				PD comorbid			
	B	OR	95%CI	p	B	OR	95%CI	p	B	OR	95%CI	p
Somatic Trait Anxiety	0.18	1.20	1.16–1.25	0.0001	0.19	1.20	1.15–1.26	0.000	0.17	1.19	1.14–1.25	0.000
Psychic Trait Anxiety	0.02	1.02	0.97–1.07	0.419	–0.04	0.97	0.92–1.01	0.160	0.05	1.05	0.98–1.12	0.143
Stress Susceptibility	0.05	1.05	1.01–1.10	0.013	0.03	1.03	0.98–1.08	0.296	0.06	1.06	1.01–1.12	0.022
Lack of Assertiveness	–0.06	0.94	0.90–0.98	0.003					–0.06	0.94	0.90–0.99	0.014
Impulsiveness	–0.01	0.99	0.95–1.03	0.597					–0.03	0.98	0.93–1.02	0.257
Adventure Seeking	–0.04	0.96	0.93–0.99	0.021								
Detachment									–0.02	0.98	0.94–1.03	0.484
Social Desirability	–0.02	0.98	0.95–1.01	0.208	–0.04	0.97	0.93–1.00	0.081	–0.02	0.98	0.94–1.03	0.399
Embitterment	–0.04	0.96	0.93–1.00	0.069					–0.01	0.99	0.94–1.04	0.686
Trait Irritability	0.03	1.03	0.99–1.09	0.183	0.00	1.00	0.95–1.04	0.867	0.05	1.05	0.99–1.12	0.093
Mistrust	–0.02	0.98	0.95–1.02	0.292					–0.04	0.96	0.92–1.00	0.070
Verbal Trait Aggression	–0.00	0.99	0.96–1.04	0.95					0.00	1.00	0.95–1.06	0.943
Physical Trait Aggression									–0.03	0.97	0.93–1.02	0.204

OR= odds ratio, 95%CI= 95% confidence intervals

6.2.3. Personality traits in patients with pure and comorbid PD

Comparison of comorbid PD and healthy controls showed significant differences in all SSP traits, except of Adventure Seeking; all traits, apart from Social Desirability with lower scores, were more expressed in comorbid PD group. Neuroticism and Aggressiveness, but not Extraversion showed significant differences between the groups (Table 5). Multivariate regression analyses in PD comorbid group showed significant differences remaining for Somatic Trait Anxiety, Stress Susceptibility and Lack of Assertiveness (Table 6). Patients with pure PD were characterized by increased anxiety-related traits, Stress Susceptibility, Irritability, and lower Social Desirability, whereas the scores for other traits did not significantly differ from those in controls. Additionally, the patients with PD only, demonstrated higher Neuroticism and Aggressiveness, but similar Extraversion, as compared to healthy subjects (Table 5). In multivariate analyses only Somatic Trait Anxiety remained significant for the group of PD only (Table 6). The comparison of the two PD groups showed that Anxiety Traits, Stress Susceptibility, Embitterment and Irritability were significantly more pronounced in patients with comorbid PD than in those with PD only, with no significant differences in other SSP traits. In respect to SSP factors, only Neuroticism significantly differed between these two PD groups (Table 5).

6.2.4. Associations of clinical and demographic factors with the personality

The patients with current PD had significantly higher scores for anxiety traits and Stress Susceptibility, but not for any other SSP traits, than PD patients in remission. Accordingly, only Neuroticism was significantly higher in patients with current PD ($p=0.0002$) as compared to the remitters. Patients with current PD showed significant differences from healthy controls on several SSP subscales, except Impulsiveness, Detachment and Physical Trait Aggression. On the other hand, patients with PD in remission reported six SSP subscales as similar to controls, including the same three as the patients with current PD, plus Lack of Assertiveness, Adventure Seeking and Mistrust. Nevertheless, both remitters and currently symptomatic patients demonstrated significantly higher Neuroticism and Aggressiveness as compared to healthy subjects (data are not shown). Comparisons of patients with earlier and later onset of PD or patients with and without family history of psychiatric disorders showed no significant differences on any SSP subscales or factors (data are not shown). Only Trait Irritability was significantly higher in PD with agoraphobia as compared to PD without agoraphobia ($p=0.0028$). In pure PD group no significant differences in SSP traits emerged between patients with and without agoraphobia. Although all three SSP factors showed higher scores in PD with agoraphobia, only Aggressiveness withstood Bonferroni correction ($p=0.02$ for Neuroticism, $p=0.04$ for Extraversion and $p=0.0028$ for Aggressiveness). SSP scores in PD group were not associated with gender, somatic morbidity, marital or occupational status (data are not shown).

6.2.5. Correlation data

The correlations between individual SSP traits and four variables, including age, onset of PD, education and number of smoked cigarettes per day, were examined in PD group. The analyses revealed modest, but significant correlations between age of patients and social desirability ($r=0.18$; $p=0.01$), onset of PD and Verbal Trait Aggression ($r=-0.19$; $p=0.01$), educational level and Embitterment ($r=-0.20$; $p=0.006$) and Mistrust ($r=-0.21$; $p=0.005$), and between smoking and Somatic Trait Anxiety ($r=0.16$; $p=0.03$) and Physical Trait Aggression ($r=0.17$; $p=0.025$). In addition, several correlations were observed for SSP factors (Table 4). In particular, Neuroticism was negatively correlated with years of education in PD group and in all studied subjects. In all subjects a positive correlation was seen between Neuroticism and numbers of smoked cigarettes per day. Extraversion was negatively correlated with age in control and total group, but not in PD sample. There were significant correlations between Aggressiveness and age as well as onset of PD and number of cigarettes in PD group. Aggressiveness was negatively correlated with age in both control and total group as well as positively correlated with number of smoked cigarettes in total sample of subjects (Table 7).

Table 7. Correlation analyses of SSP factors in patient and control groups

	Neuroticism	Extraversion	Aggressiveness
PD group (n=193)			
Age (years)	$r=-0.32$; $p=0.66$	$r=0.01$; $p=0.86$	$r=-0.16$; $p=0.03$
Onset of PD (years)	$r=-0.03$; $p=0.73$	$r=-0.04$; $p=0.60$	$r=-0.15$; $p=0.04$
Education (years)	$r=0.16$; $p=0.03$	$r=0.01$; $p=0.91$	$r=-0.03$; $p=0.72$
Cigarettes (N)	$r=0.09$; $p=0.24$	$r=0.02$; $p=0.81$	$r=0.18$; $p=0.01$
Control group (n=314)			
Age (years)	$r=0.05$; $p=0.43$	$r=-0.18$; $p=0.001$	$r=-0.23$; $p=0.000$
Education (years)	$r=0.10$; $p=0.08$	$r=0.02$; $p=0.79$	$r=-0.08$; $p=0.17$
Cigarettes (N)	$r=0.01$; $p=0.80$	$r=-0.01$; $p=0.86$	$r=0.11$; $p=0.06$
All subjects (n=507)			
Age (years)	$r=-0.03$; $p=0.46$	$r=-0.11$; $p=0.01$	$r=-0.21$; $p=0.000$
Education (years)	$r=-0.13$; $p=0.004$	$r=0.01$; $p=0.77$	$r=-0.07$; $p=0.11$
Cigarettes (N)	$r=0.10$; $p=0.02$	$r=0.00$; $p=0.97$	$r=0.17$; $p=0.000$

6.2.6. Discussion

We presented the first study using SSP to show significant differences in the self-reported personality traits of patients with PD as compared to healthy subjects. Earlier, higher Neuroticism and probably lower Extraversion were found in patients with PD who were assessed with NEO or Maudsley Personality Inventory scales, whereas other personality traits were reported to be

similar to those in healthy subjects (Bienvenu et al., 2001a, Bienvenu et al., 2004, Carrera et al., 2006, Freire et al., 2007). As a personality pattern of PD patients has been characterised by higher rates of pathological personality traits compared to controls, our results showed a deviation of practically all SSP traits in patients with PD. We demonstrated that Neuroticism was the main personality characteristic of patients with PD, which was in a line with previous reports (Bienvenu et al., 2005, Freire et al., 2007, Zugliani et al., 2017). In particular, increased Somatic and Psychic Trait Anxiety and Stress Susceptibility were observed in all PD groups regardless of clinical status or phenotypes. By considering all personality traits simultaneously, Somatic Trait Anxiety stood out as the strongest predictor of the presence of PD. The SSP Somatic Trait Anxiety indicates a tendency to experience autonomic arousal; therefore, our findings are in line with the later developments of the tripartite model showing a specific link between heightened arousal and PD (Brown et al., 1998, Clark et al., 1994) as well as with the cognitive model of PD, which emphasises the role of increased interoceptive sensitivity and misinterpretation of somatic symptoms (Clark, 1986) along with the corresponding importance of anxiety sensitivity as a trait underlying the tendency for catastrophic misinterpretations of bodily changes (McNally, 2002).

Notably, the degree of Neuroticism correlated negatively with education level in both the PD group and the total sample of studied subjects, indicating that lower education may contribute to higher neurotic behaviour. One of the new findings was the significantly higher Aggressiveness among PD patients as compared to healthy subjects. Furthermore, the increased Aggressiveness was seen in both pure and comorbid PD groups, but it was more pronounced in the comorbid group, suggesting that aggression-related traits in PD may be associated with affective comorbidity. Another factor related to Aggressiveness in our sample was the presence of agoraphobia. Interestingly, PD patients with agoraphobia showed significantly higher Aggressiveness and particularly increased Trait Irritability than patients without agoraphobia. This finding suggests a link between the psychopathology of agoraphobia and trait Aggressiveness, which requires further research. In contrast to some of the previous studies, we found no difference in Extraversion between any of the studied PD groups and controls, suggesting that this personality aspect is probably less affected in PD.

Previous studies have proposed an accentuation of some personality traits, including Harm Avoidance, Neuroticism and Extraversion, in PD patients with affective comorbidity as compared with those experiencing pure PD (Ampollini et al., 1997, Bienvenu et al., 2001b, Freire et al., 2007). Similarly, Cuijpers et al. (2005) found Neuroticism to be the trait most strongly related to comorbidity. In our study, practically all SSP subscale scores were higher in patients with comorbid compared to pure PD. However, among SSP factors, only Neuroticism was higher among patients with comorbid panic phenotype. A number of clinical and demographic variables in our study were similar between PD only and comorbid groups, except the higher number of subjects with family psychiatric history among patients with affective comorbidity. Nevertheless,

none of the SSP traits or factors was significantly related to family psychiatric history. Thus, it is unlikely that increased Neuroticism in the comorbid PD group could be explained by these factors.

We also examined the relationships of personality traits to several demographic and clinical characteristics. The patients with current PD showed higher Neuroticism and related traits than PD patients in remission, but they did not differ in any other SSP trait or factor. Bienvenu et al. (2004) had also detected higher Neuroticism in patients with current PD compared to those in remission. Although high Neuroticism was proposed to predict the later onset of anxiety disorders (Bienvenu, 2007), none of the SSP personality traits significantly differed between patients with the earlier (<30 years) and late (>29 years) onset of PD in our sample. Previous studies have found that low Extraversion and low trust were associated with both agoraphobia and social phobia, but not with depression (Bienvenu et al., 2001a, Bienvenu et al., 2001b, Bienvenu et al., 2004). Furthermore, Carrera et al. (2006) have shown that PD patients were only more introverted than controls in the presence of agoraphobia. However, higher Neuroticism in patients with PD was prominent irrespective of agoraphobia. Furthermore, Openness, Agreeableness and Conscientiousness were unrelated to agoraphobia in their study. In our data set, neither Extraversion nor Neuroticism was related to agoraphobia. In respect to gender effects, Foot and Koszycki (2004), using NEO-PI-R, showed a lack of significant differences in anxiety sensitivity and Neuroticism scores between female and male PD patients with or without agoraphobia. However, women scored higher than men on Extraversion and men on angry hostility and depression facets of Neuroticism. In our study, none of the SSP traits was significantly affected by gender.

To the best of our knowledge, the sample of patients with PD in our study was to date the largest among the studies investigating personality traits in PD. However, certain limitations should be considered when interpreting our results. We were unable to account for the effects of treatment on personality traits in our sample. Notably, neither clinical severity nor short-term treatment response in PD was influenced by personality factors in an earlier study by Carrera et al. (2006). Both our findings and previous findings indicate that the expression of anxiety and stress-related traits may be state-dependent and intensified in the symptomatic stage of PD. We did not use specific clinical scales for the assessment of anxiety and panic severity. Although current PD would evidently be associated with more severe symptoms than PD in remission, the correlations between clinical and personality assessments should be addressed in more detail in the future. A possible confounder in our study was the small proportion of patients who met diagnostic criteria for bipolar disorder or (hypo)manic episode. Although this subgroup demonstrated personality traits consistent with non-bipolar groups (data not shown), the specific characteristics of PD with bipolar comorbidity need to be investigated in a larger sample.

6.3. Study III: Association between personality traits and escitalopram treatment efficacy in PD

6.3.1. Demographic and clinical characteristics

In present study, out of the 107 patients, who completed the study, 38 were diagnosed with PD (36%), 24 with PD and agoraphobia (22%), 14 with PD and co-morbid mild depression (13%) and 31 with PD, agoraphobia and mild depression (29%). The demographic characteristics of these groups did not differ significantly (Table 8). All 107 patients had a PDSS score higher than 7. At the beginning of the study, the mean score of PDSS was 16.07 and by the end of the study 5.16. By the end of study eighty-four (79%) patients met the criteria of remission.

Table 8. Demographic and clinical data

Variable	Mean or Frequency (n)	SD or %
Age (years; mean, sd)	34	11,73
Education (years; mean, sd)	13,7	2,5
Duration of PD (years; mean, sd)	7,7	9,5
Comorbid somatic diseases (n, %)	37	26
Gender (n, %)		
Male	36	34
Female	71	66
Occupation (n, %)		
Employed	70	67
Student	14	13
Retired	2	2
Unemployed	17	16
Disability	1	1
Smoking (n, %)		
Never	54	52
Formerly	22	21
Currently	28	27
Alcohol (n, %)		
Does not use	22	21
Once a month	39	38
2–4 times a month	35	34
2–3 times a week	8	8
Diagnoses (n, %)		
PD	38	36
PD and agoraphobia	24	22
PD and mild depression	14	13
PD, agoraphobia, and mild depression	31	29

6.3.2. Changes in SSP sub-scale scores following escitalopram treatment in PD

The means of all the T-scores of SSP sub-scales express tendency to decrease during the study. The largest changes were in Somatic Trait Anxiety (-6.59 ± 9.98), Stress Susceptibility (-5.22 ± 9.55), and Psychic Trait Anxiety (-5.13 ± 8.18). The rest of the means of the scores of SSP sub-scales decreased less (Table 9).

Table 9. SSP sub-scale scores at visit 0 and visit 6, and the change of the scores

SSP sub-scale	Baseline; mean (sd) min...max	Study end; mean (sd) min...max	Difference; mean (sd) min...max	<i>p</i> -value for difference
Somatic Trait Anxiety	65.91 (9.14) 45.02...89.41	59.32 (10.62) 34.06...86.8	-6.59 (9.98) -28.72...18.28	<0.0001*
Psychic Trait Anxiety	60.27 (9.67) 33.89...79.51	55.13 (9.55) 33.89...72.31	-5.13 (8.18) -31.21...24.01	<0.0001*
Stress Susceptibility	60.3 (10.59) 34.12...86.71	55.08 (11.06) 28.28...80.86	-5.22 (9.55) -26.29...37.98	<0.0001*
Lack of Assertiveness	52.43 (10.95) 23.43...77.58	49.52 (10.1) 26.4...80.57	-2.9 (8.48) -41.58...17.9	0.0009*
Impulsiveness	46.69 (11.51) 16.69...84.38	44.58 (11.38) 16.69...75.04	-2.11 (7.94) -26.83...18.67	0.0059*
Adventure Seeking	46.52 (10.13) 21.62...67.46	45.66 (10.18) 24.17...72.55	-0.86 (6.46) -20.81...13.01	0.3531
Detachment	50.16 (9.83) 24.05...73.69	49.45 (10.12) 23.68...70.57	-0.71 (6.88) -19.62...22.88	0.2695
Social Desirability	48.04 (10.35) 23.12...72.26	46.54 (10.51) 23.12...72.26	-1.51 (7.72) -28.08...14.04	0.0834
Embitterment	55.99 (10.48) 32.51...91.78	53.16 (11.12) 32.51...82.89	-2.82 (7.3) -23.72...11.86	0.0004*
Trait Irritability	57.17 (9.08) 32...77.35	53.91 (9.93) 26.78...75.08	-3.26 (7.08) -24.95...11.33	<0.0001*
Mistrust	53.62 (11.27) 28.14...89.21	51.68 (11.59) 25.22...83.44	-1.93 (9.22) -34.63...29.15	0.0240
Verbal Trait Aggression	52.52 (9.54) 28.55...75.99	51.09(9.7) 31.04...75.99	-1.42 (7.93) -44.88...14.96	0.1668
Physical Trait Aggression	52.37 (10.42) 33.11...84.02	51.77 (11.19) 30.72...86.54	-0.6 (6.85) -25.2...15.12	0.7796

* significant after Holm-Bonferroni correction for multiple comparisons

6.3.3. Correlation data

None of the baseline scores of SSP sub-scales was strongly associated with the treatment response. Only the Impulsiveness score was weakly correlated with the change of PDSS ($r=0.22$, $p=0.033$), and it helped to predict the odds for remission OR=0.90, $p=0.005$); however, these findings were statistically not significant after the adjustment for multiple comparisons (Table 10).

Table 10. Correlations between baseline scores of SSP sub-scales and change in PDSS and ratios of the odds of remission

SSP sub-scale	r (95% CI)	P	OR* (95% CI)	p
Somatic Trait Anxiety	0.05 (−0.16...0.24)	0.675	1.02 (0.93...1.03)	0.697
Psychic Trait Anxiety	−0.04 (−0.23...0.18)	0.818	1.00 (0.95...1.05)	0.928
Stress Susceptibility	0.04 (−0.16...0.24)	0.692	0.98 (0.96...1.05)	0.623
Lack of Assertiveness	0.01 (−0.18...0.23)	0.810	1.05 (0.95...1.04)	0.240
Impulsiveness	0.22 (0.03...0.41)	0.033	0.90 (0.88...0.97)	0.005
Adventure Seeking	−0.01 (−0.22...0.19)	0.895	1.02 (0.97...1.06)	0.590
Detachment	−0.03 (−0.24...0.16)	0.692	1.02 (0.97...1.06)	0.572
Social Desirability	−0.05 (−0.23...0.17)	0.766	1.04 (0.97...1.07)	0.254
Embitterment	0.00 (−0.18...0.23)	0.818	0.96 (0.91...1.00)	0.239
Trait Irritability	0.13 (−0.05...0.35)	0.142	0.94 (0.90...1.00)	0.152
Mistrust	0.09 (−0.11...0.29)	0.401	0.98 (0.91...0.99)	0.521
Verbal Trait Aggression	0.09 (−0.07...0.32)	0.222	1.00 (0.94...1.04)	0.898
Physical Trait Aggression	0.20 (−0.01...0.38)	0.064	1.00 (0.97...1.06)	0.999

* Adjusted for sex, age, years of education, level of drinking frequency, presence of somatic comorbidities, and for the presence of mild depression and/or agoraphobia. The remission was defined if the score on the CGI improvement scale was 2 or less, the PDSS score 7 or less and no panic attacks had occurred for at least last two weeks.

Some weak correlations existed between the changes of the scores of SSP sub-scales and the changes in the clinical assessment scores. The change of SSP Stress Susceptibility was weakly correlated with the change of CGI-S ($r=-0.27$); the change of SSP Lack of Assertiveness was weakly correlated with the change of HAM-A ($r=0.32$) and EST-Q depression ($r=0.21$). However, none of the partial correlation coefficients was statistically significant after Holm-Bonferroni correction. (Table 11). The baseline scores of SSP sub-scales and the change in EST-Q depression score were not correlated. Similarly, neither changes in CGI-S scores nor changes in HAM-A scores were correlated with the baseline scores of SSP sub-scales. Partial correlations between Impulsiveness and change in CGI-S ($r=0.29$) and change in HAM-A ($r=0.28$) were statistically not significant after correcting for multiple testing (Table 12).

Table 11. Correlations between the changes of scores of SSP sub-scales and the changes of clinical assessment scores

	PDSS*	CGI-S*	HAM-A*	EST-Q depression*
Somatic Trait Anxiety	0.07 (-0.13...0.27)	-0.16 (-0.35...0.05)	0.24 (0.04...0.43)	0.03 (-0.18...0.23)
Psychic Trait Anxiety	0.11 (-0.09...0.31)	-0.09 (-0.29...0.12)	0.15 (-0.06...0.34)	0.10 (-0.10...0.30)
Stress Susceptibility	0.15 (-0.06...0.34)	-0.27 (-0.45...-0.07)	0.20 (0.00...0.39)	0.10 (-0.10...0.30)
Lack of Assertiveness	0.07 (-0.14...0.27)	-0.12 (-0.32...0.08)	0.32 (0.13...0.50)	0.21 (0.01...0.39)
Impulsiveness	-0.08 (-0.28...0.12)	0.07 (-0.14...0.27)	-0.07 (-0.27...0.14)	-0.02 (-0.22...0.18)
Adventure seeking	0.09 (-0.12...0.28)	-0.07 (-0.27...0.14)	0.09 (-0.12...0.29)	0.10 (-0.11...0.29)
Detachment	0.09 (-0.11...0.29)	0.02 (-0.18...0.23)	-0.01 (-0.22...0.19)	-0.09 (-0.28...0.12)
Social Desirability	-0.03 (-0.23...0.17)	0.09 (-0.12...0.29)	-0.02 (-0.22...0.19)	0.02 (-0.18...0.22)
Embitterment	0.16 (-0.04...0.35)	-0.17 (-0.36...0.04)	-0.01 (-0.21...0.19)	0.00 (-0.20...0.20)
Trait Irritability	-0.01 (-0.21...0.19)	0.06 (-0.15...0.26)	0.06 (-0.14...0.26)	0.04 (-0.17...0.23)
Mistrust	-0.05 (-0.25...0.16)	-0.02 (-0.22...0.19)	-0.07 (-0.27...0.14)	0.10 (-0.10...0.30)
Verbal Trait Aggression	-0.14 (-0.33...0.07)	0.07 (-0.13...0.27)	-0.06 (-0.26...0.15)	-0.15 (-0.34...0.05)
Physical Trait Aggression	-0.09 (-0.29...0.11)	0.03 (-0.17...0.24)	-0.13 (-0.32...0.08)	-0.10 (-0.29...0.11)

* *p*-values were statistically significant after Holm-Bonferroni correction for multiple comparisons. Adjusted for sex, age, years of education, level of drinking frequency, presence of somatic co-morbidities, and for the presence of mild depression and/or agoraphobia.

Table 12. Correlations between baseline scores of SSP sub-scales and changes in CGI-S, HAM-A, and EST-Q depression

	CGI-S		HAM-A		EST-Q depression	
	<i>r</i> (95% CI)	<i>p</i> *	<i>r</i> (95%CI)	<i>p</i> *	<i>r</i> (95% CI)	<i>p</i> *
Somatic Trait Anxiety	-0.09 (-0.29...0.12)	0.407	-0.06 (-0.26...0.15)	0.608	0.03 (-0.18...0.22)	0.817
Psychic Trait Anxiety	-0.03 (-0.23...0.18)	0.806	0.04 (-0.17...0.24)	0.713	0.15 (-0.05...0.34)	0.152
Stress Susceptibility	-0.04 (-0.24...0.17)	0.740	0.11 (-0.10...0.30)	0.329	0.20 (-0.01...0.38)	0.063
Lack of Assertiveness	-0.08 (-0.28...0.13)	0.486	-0.08 (-0.28...0.13)	0.476	0.12 (-0.09...0.31)	0.278
Impulsiveness	0.30 (0.10...0.47)	0.005	0.29 (0.09...0.46)	0.007	0.06 (-0.15...0.26)	0.589
Adventure Seeking	0.04 (-0.17...0.24)	0.723	-0.04 (-0.24...0.17)	0.738	-0.13 (-0.33...0.07)	0.206
Detachment	-0.07 (-0.28...0.13)	0.493	0.03 (-0.17...0.24)	0.754	0.13 (-0.08...0.32)	0.234
Social Desirability	0.04 (-0.17...0.24)	0.744	-0.01 (-0.22...0.19)	0.915	-0.10 (-0.30...0.10)	0.326
Embitterment	-0.02 (-0.23...0.18)	0.838	0.09 (-0.11...0.29)	0.394	0.10 (-0.10...0.30)	0.342
Trait Irritability	0.18 (-0.03...0.37)	0.098	0.20 (0.00...0.39)	0.061	0.04 (-0.16...0.24)	0.712
Mistrust	-0.04 (-0.24...0.17)	0.727	0.23 (0.03...0.41)	0.032	0.09 (-0.11...0.28)	0.406
Verbal Trait Aggression	0.12 (-0.08...0.32)	0.256	0.14 (-0.07...0.33)	0.210	0.09 (-0.12...0.28)	0.408
Physical Trait Aggression	0.14 (-0.06...0.34)	0.186	0.28 (0.08...0.46)	0.008	0.03 (-0.17...0.23)	0.794

* uncorrected *p*-values; all were statistically not significant after Holm-Bonferroni correction for multiple comparisons; 95% confidence intervals and *p*-values, adjusted for sex, age, years of education, level of drinking frequency, presence of somatic co-morbidities, and for the presence of mild depression and/or agoraphobia.

6.3.4. Discussion

Several personality traits on the SSP measurement, including Embitterment, Impulsiveness, Lack of Assertiveness, Psychic and Somatic Trait Anxiety, and Stress Susceptibility, improved after 12 weeks of medication in comparison with the baseline scores. Although the changes in the SSP scores after the treatment period were generally more pronounced among remitters to escitalopram treatment compared to non-remitters, none of scores of the SSP significantly differed between these groups at the end of the treatment period. Moreover, no significant correlations were found between the changes of scores of SSP subscales and the changes of clinical assessment scores following 12 weeks of medication. These results look quite intriguing in the light of our previous study, which demonstrated the maladaptive personality disposition of practically all SSP traits in patients with PD (Võhma et al., 2010, Study II), confirming that maladaptive trait patterns may contribute to the development and expressions of PD. The current study suggests that antidepressant medication is accompanied by certain improvement in personality patterns in patients with PD, but changes in the traits do not seem to be clearly associated with treatment efficacy. Although our results may confirm the trait rather than state phenomena of personality deviations in PD, it is possible to speculate that personality characteristics are more flexible in terms of normalisation, at least among good responders to pharmacological intervention. The effect of antidepressant medication on personality traits in PD has not been univocally proven. None of the personality traits assessed by the Neuroticism-Extraversion-Openness Five-factor Personality Inventory was related to the short-term response to SSRI treatment in PD (Carrera et al., 2006). Still, a significant decrease was observed in all anxiety related traits after 6 months of treatment with SSRI citalopram in a smaller sample of PD patients, using the original scale for SSP measurement, the KSP (Neuger et al., 2002). Considering that a similar tendency to decrease SSP scores was found among our patients, we may suggest that a longer treatment period would probably reveal a statistically stronger effect of medication on personality traits. However, the more pragmatic explanation would be that the response to antidepressant medication and particularly the substantial decrease in clinical severity could also improve the self-rating of one's own personality characteristics.

The second main goal of our study was to detect the possible predictive effect of personality traits on treatment response to antidepressant medication. The only positive finding we detected, but which did not withstand Bonferroni correction for multiple comparisons, was the predictive effect of impulsivity. PD non-remitters to 12-week treatment with escitalopram showed a higher Impulsivity score at baseline SSP assessment compared with responders. SSP Impulsivity correlates with several constructs of the five-factor personality model. The strongest associations have been found with the Impulsiveness facet of Neuroticism and the Deliberation facet of Conscientiousness, thereby combining a low frustration tolerance with non-planning impulsivity (Aluoja et al., 2009).

Lower Neuroticism has predicted clinical improvement at three-month point in a Collaborative Care for Anxiety and Panic Study study (Chavira et al., 2009). The predictive effect of Impulsivity in our study was also confirmed by correlation analyses between SSP and the clinical scales scores. In particular, the higher score of Impulsivity at baseline was close-to-significantly associated with a lesser decrease in the CGI severity score during the treatment period. The same trends were also seen on the PDSS and HAM-A scales, but correlations did not reach statistically significant levels. Previously we reported that PD phenotype is characterised by a higher Impulsivity in comparison to normal presentation (Võhma et al., 2010, publication II). Nevertheless, the relationship between anxiety and impulsivity seems controversial (Barratt 1965; Askenazy et al., 2000). Our study accords with Marchesi et al. (2006) in finding that features of borderline personality disorder predicted a lower likelihood of remission with SSRIs in PD patients. Emotional instability and low impulse control are characteristic features of borderline personality, and they are also part of the Impulsivity construct measured by the SSP. Studies conducted with different methodologies have found no correlation between anxiety and impulsivity (Askenazy et al., 2000; Apter et al., 1993; Lecrubier et al., 1995; Caci et al., 1998). On the other hand, in several studies anxiety disorder patients have reported higher scores of impulsivity than healthy controls (Summerfeldt et al., 2004; Del Carlo et al., 2012). The possible involvement of impulsivity in treatment response appears to be less understood. Silva et al. (2010) demonstrated that borderline personality disorder carriers of long alleles of serotonin transporter promoter region related polymorphism (5-HTTLPR) had a better anti-impulsive response to SSRI fluoxetine treatment than in patients with short alleles. Walderhaug et al. (2010) also showed that reduced serotonergic neurotransmission following acute tryptophan depletion increased impulsivity in healthy men, but decreased it in women. Considering that the relationship between reduced function of the central serotonin system and impulsivity is well recognised (Brown et al., 1979; Brown et al., 1982; Coccaro et al., 1989; O'Keane et al., 1992), we suggest that an increased impulsivity and worse anti-panic response to escitalopram treatment may indicate a more pronounced or persistent serotonergic dysfunction, at least in the sub-group of patients with poor treatment outcome. Interestingly, significant and persistent changes in central serotonin functioning were observed by neuroimaging studies in male patients with PD (Maron et al., 2010b; Nash et al., 2008; Cannon et al., 2013). Nevertheless, none of these brain-imaging studies had measured the expression of the Impulsivity trait among their sample.

To the best of our knowledge, the sample of patients with PD in our study was to date the largest among the studies investigating the relationship between personality traits and treatment response in PD. However, certain limitations should be taken into account when interpreting our results. All patients were treated in a naturalistic way using an open-label drug, so placebo responses cannot be excluded; the control group is absent. However, to diminish possible sources of bias in variation in clinical assessments, the severity of PD symptoms

and treatment response were carefully rated by one experienced psychiatrist at each visit. Furthermore, all clinical assessments on the patients' self-reports on SSP measurements were made blindly. However, our sample size can still be considered not large enough to reveal statistically significant findings, especially keeping in mind several other factors, such as gender, age and co-morbidity.

7. GENERAL DISSCUSSION

According to our results, the Estonian version of the SSP has demonstrated high reliability and validity as evaluated in reference to both: the Swedish original data and the position of the SSP-measured traits within the basic personality dimensions of the five-factor model. Since the first introduction of SSP (Gustavsson et al., 2000), this scale has been applied in various mental health and somatic disorders, including depression, schizophrenia, bipolar disorder, cognitive disturbances, eating problems, psoriasis and others (Aluoa et al., 2018; Iliadis et al., 2015; Ausén et al., 2011, Elfhag 2005; Fagerberg et al., 2018; Rasul et al., 2016; Remröd et al., 2015; Sparding et al., 2017; Volgsten et al., 2010). The initial idea of developing KSP and its later modification with improved psychometric qualities, SSP, was to attain an instrument for evaluating personality traits in psychopathological groups. This was also the main reason for adaption of the Estonian version of SSP for several purposes, including the personality assessment in research and a clinical context. The SSP appears to be fairly stable in terms of repeated measures within the same individual groups (Hedman et al., 2014, Muller et al., 2015), and has shown similar factor loadings in different samples (Gustavsson et al., 2000, Fagerber et al., 2016, Volgsten H et al., 2010). The other advantage is its relatively short format, consisting of 91 items, as compared to 238–240 items for the full NEO and TCI questionnaires, which is an argument in favour of time effectiveness for SSP. Thus in line with earlier studies, the current work gives additional strong support for recommending the extensive use of SSP in further exploration of the role of personality factors in mental disorders.

It should be noted that the current work is the first implementation of SSP in investigation of personality traits in PD, which is very important in that the relationship between personality traits and PD is significantly less explored than the association of PD with personality disorders. We found that virtually all SSP traits show an abnormal range in patients with PD, while at least some personality aberrations seem to be more pronounced in PD with affective comorbidity compared to pure PD. Generally, the SSP factors of Neuroticism and Aggressiveness, but not Extraversion, were significantly higher in the PD group than in controls, and only a few demographic and clinical variables were associated with SSP scores in PD patients. These results add to the evidence of a maladaptive personality disposition in patients with PD, particularly high Neuroticism and manifest Somatic Trait Anxiety, which are probably not influenced by other clinically relevant factors. The associations between PD and both higher Neuroticism as well as lower Extraversion are in line with similar earlier observations (Bienvenu et al., 2001b, Bienvenu et al., 2004, Carrera et al., 2006, Freire et al., 2007). However, the finding of higher Aggressiveness and increased Trait irritability among PD patients, particularly with agoraphobia, as compared to healthy subjects seems very intriguing. In our previous study, we examined how personality disposition as assessed with SSP may affect the

response to the cholecystokinin tetrapeptide (CCK-4; 50 microg) challenge in healthy volunteers (n=105). Our results showed that the occurrence of CCK-4-induced panic attacks was best predicted by adhering to SSP-defined traits: Lack of Assertiveness, Detachment, Embitterment and Verbal Aggression. Moreover, for different subsets of CCK-4-induced symptoms, the traits of Physical Aggression, Irritability, Somatic Anxiety and Stress Susceptibility also appeared related to panic manifestations. In particular, higher Verbal trait Aggression predicted the occurrence of a panic attack, whereas increased Physical Trait Aggression predicted all aspects of panic symptoms (Toru et al., 2010). Previously, it has also been indicated that the tendency to be verbally aggressive was associated with a panic response to pentagastrin administration (Radu et al., 2003). The association between Aggressiveness and provoked panic response is further supported by George et al. (2000), who showed that persons exhibiting a high rate of physical aggression in domestic relationships react with more panic symptoms to sodium lactate provocation. The exact nature of these associations is unclear. A Radu et al. (2003) finding that personality emerges as a significant predictor of panic reaction only with a lower dose of pentagastrin suggests a rather non-specific stress reactivity. It is possible that certain personality traits heighten an individual's reactivity to different kinds of stressors, including panic-provoking agents. On the other hand, anxiety and aggression are known to share underlying neurobiology, including aspects of CCK neurotransmission and pharmacotherapy approaches (Siegel et al., 2007). Thus, our observation that Aggressiveness predisposes to a greater panic reaction in a laboratory challenge in healthy volunteers may indicate one relationship between traits of aggression and susceptibility to panic attacks. However, the relevance of these findings to clinical aspects of PD is not fully clear. Although an earlier study found that patients with PD reported significantly greater levels of anger aggression compared to controls (Moscovitch et al., 2008), this was not confirmed by other research showing a non-significant relationship between PD and aggression (Niazi et al., 2008). Interestingly, neither Physical nor Verbal traits of Aggression changed throughout the 12-week treatment with escitalopram in the current study, suggesting that aggressiveness is probably one of the endophenotypic features of PD.

Although several other personality traits, including Embitterment, Impulsiveness, Lack of Assertiveness, Psychic and Somatic Trait Anxiety and Stress Susceptibility, have improved on the SSP measurement after 12 weeks treatment with escitalopram, none of these positive changes significantly differed between remitters and non-remitters at the end of the treatment period. Furthermore, the lack of correlations between the changes in personality traits and the changes of clinical assessment scores following escitalopram treatment indicates that improvement in personality characteristics is probably contributed by the general effect of recovery resulting in better self-perception. Considering the earlier suggestion that the result of therapy can be attributed to the state level variance in personality trait measures (Du et al., 2002; Marchevsky, 1999), it would be tentative to speculate that SSP improvement following a treatment

period in PD can be temporally and may depend on the disease course. However, the recent meta-analysis conducted by Roberts et al. (2017) based on data from published studies in 1959–2013 and involving over 20,000 participants demonstrated that in comparison with other disorders such as depression, eating disorders, personality disorders and substance use, patients with anxiety disorders reported more pronounced and long-lasting changes in personality characteristics, especially in Neuroticism and Extraversion, just after a few weeks of therapy. In addition, a brief internet-based cognitive behaviour therapy (12 weeks) for severe health anxiety causes long-term changes in measures of SSP personality traits related to Neuroticism, whereas Extraversion and Aggression remained largely unchanged (Hedman et al., 2014). Interestingly, the stability of the SSP personality trait was also reasonably high among patients with psychotic disorders, although lower than among non-psychotic individuals in the study of Fagerberg (2018). On the other hand, Roberts et al. (2017) concluded by meta-analysis of earlier studies that psychopharmacological therapies actually exhibited a slightly smaller effect than the other treatment approaches, e.g. CBT, supportive therapy and psychodynamic methods, on changes of personality traits. This observation looks very intriguing in terms of further exploration of non-pharmacological therapy methods and their influence on SSP personality traits in PD patients.

The other dilemma raised from this study is the predictive effect of personality traits on treatment response to antidepressant medication in PD. The only personality trait that shows the possible predictive effect in current study was Impulsivity. In particular, a higher Impulsivity score at baseline SSP measurement in PD patients was associated with poor treatment outcome; however, no other SSP personality characteristics demonstrated a predictive effect. These results are in accordance with our recent study aiming to identify specific personality traits that could predict the treatment response and/or the dynamics of symptom change in 132 outpatients with major depressive disorders, who were also treated with escitalopram antidepressants for 12 weeks. We found that none of the studied SSP traits predicted the end result of the treatment. However, patients with higher Social Desirability, lower Stress Susceptibility and lower Irritability achieved a more rapid decrease in depression symptoms. Nevertheless, the latter associations did not survive the multiple testing corrections, probably due to the relatively small sample (Aluoja et al., 2018). Therefore, these findings suggest that specific personality traits may predict the trajectory of symptom change rather than the overall improvement rate, which is probably true for both highly co-existing mental disorders such as PD and depression. Notably, the lack of the major influence of SSP personality on the disease course was reported in patients with bipolar disorder, indicating that the personality profile does not seem to have prognostic value over a 2-year period in this type of mood disorder (Sparding et al., 2017).

Finally, it should be noted that genetic factors, not investigated in this study, should be considered when exploring the relationship between PD and personality traits. For example, there are recent findings showing that genetic poly-

morphisms, particularly within hydroxysteroid (11-beta) dehydrogenase 1 gene and the serotonin transporter gene-linked polymorphism 5-HTTLPR, may modify manifestation of Neuroticism-related personality traits assessed by SSP (Iliadis et al. Gingnell et al., 2010). Although there is strong evidence that several genes may have a modulatory role on the onset, course and treatment outcome in PD, including among the Estonian population (Maron et al., 2010a), the genetic part was not included in this study due to the relatively small sample and risk of false findings; however, we recognise that the genetic impact on personality traits in PD requires further investigation.

8. CONCLUSIONS

1. We demonstrated that the SSP Estonian version has acceptable psychometric properties. The SSP measures traits that seem to be universal and are in line with the major theories of personality structure.
2. The use of SSP proved to add pertinent information on clinically relevant personality dimensions in patients with PD. We showed evidence of maladaptive trait patterns in patients with PD. In particular, these patients are characterised by high Neuroticism and higher manifestation of Somatic Trait Anxiety.
3. More pronounced personality trait deviations were observed in PD with affective comorbidity. Aggressiveness, in particular, was associated with agoraphobia and affective comorbidity.
4. Although maladaptive personality disposition in patients with PD shows a certain trend towards normalisation after a 12-week treatment with the antidepressant escitalopram, there was not a strong correlation with the clinical outcome. A better response to medication was accompanied by more pronounced improvement in personality characteristics, which is probably contributed by the general effect of recovery resulting in more positive self-perception.
5. The Impulsiveness shows a possible predictive effect on the antidepressant treatment response in PD, which seems intriguing and worthy of further investigation.

9. REFERENCES

- Akiskal HS, Hirschfeld RM, Yerevanian BI. The relationship of personality to affective disorders. *Arch Gen Psychiatry* 1983; 40 (7): 801–810.
- Almeida YA, Nardi AE. Psychological features in panic disorder: a comparison with major depression. *Arq Neuropsiquiatr* 2002; 60 (3-A): 553–557.
- Aluoja A, Shlik J, Vasar V, Luuk K, Leinsalu M. Development and psychometric properties of the Emotional State Questionnaire, a self-report questionnaire for depression and anxiety. *Nord J Psychiatry* 1999; 53: 443–449.
- Aluoja A, Tõru I, Raag M, Võhma U, Maron E. Personality traits and escitalopram treatment outcome in major depression. *Nord J Psychiatry* 2018; 72 (5): 354–360.
- Aluoja A, Voogne H, Maron E, Gustavsson JP, Võhma U, Shlik J. Personality traits measured by the Swedish universities Scales of Personality: factor structure and position within the five-factor model in an Estonian sample. *Nord J Psychiatry* 2009; 63 (3): 231–236.
- American Psychiatric Association (APA). *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV*, Fourth Edition. Washington DC: American Psychiatric Press; 1994.
- Ampollini P, Marchesi C, Signifredi R, Maggini C. Temperament and personality features in panic disorder with or without comorbid mood disorders. *Acta Psychiatr Scand* 1997; 95: 420–423.
- Andrisano C, Balzarro B, Fabbri C, Chiesa A, Serretti A. Antidepressants and Panic Disorder. *Clin Neuropsychiatry* 2012; 9 (2): 84–101.
- Apter A, Plutchik R, van Praag HM. Anxiety, impulsivity and depressed mood in relation to suicidal and violent behavior. *Acta Psychiatr Scand* 1993; 87: 1–5.
- Askenazy F, Caci H, Myquel M, Darcourt G, Lecrubier Y. Relationship between impulsivity and platelet serotonin content in adolescents. *Psychiatry Res* 2000; 94 (1): 19–28.
- Asselmann E, Stender J, Grabe HJ, König J, Schmidt CO, Hamm AO, Pané-Farré CA. Assessing the interplay of childhood adversities with more recent stressful life events and conditions in predicting panic pathology among adults from the general population. *J Affect Disord* 2018; 225: 715–722.
- Ausén B, Edman G, Almkvist O, Bogdanovic N. Self- and informant ratings of personality in mild cognitive impairment, reviewed. *Dement Geriatr Cogn Disord* 2011; 32 (6): 387–393.
- Barlow DH. *Anxiety and its disorders: The nature and treatment of anxiety and panic*. New York, NY, US: Guilford Press, 1988.
- Barlow DH, Gorman JM, Shear MD, Woods SW. Cognitive-behavioral therapy, imipramine, or their combination for panic disorder: A randomized controlled trial. *JAMA* 2000; 283 (19): 2529–2536.
- Barlow DH, Sauer-Zavala S, Carl JR, Bullis J, Ellard K. The nature, diagnosis, and treatment of neuroticism: Back to the future. *Clin Psychological Science* 2014; 2 (3): 344–365.
- Barratt ES. Factor analysis of some psychometric measures of impulsiveness and anxiety. *Psychol Rep* 1965; 16: 547–554.
- Barsky AJ, Delamater BA, Orav JE. Panic disorder patients and their medical care. *Psychosomatics* 1999; 40 (1): 50–56.

- Batelaan NM, de Graaf R, Penninx BW, van Balkom AJ, Vollebergh WA, Beekman AT. The 2-year prognosis of panic episodes in general population. *Psychol Med* 2010; 40 (1): 147–157.
- Bienvenu OJ. What is the meaning of associations between personality traits and anxiety and depressive disorders? *Braz J Psychiatry* 2007; 29: 3–4.
- Bienvenu OJ, Brandes MB. The interface of personality traits and anxiety disorders. *Primary Psychiatry* 2005; 12 (3): 35–39.
- Bienvenu OJ, Brown C, Samuels JF, Liang KY, Costa PT, Eaton WW, Nestadt G. Normal personality traits and comorbidity among phobic, panic and major depressive disorders. *Psychiatry Res.* 2001a; 102 (1): 73–85.
- Bienvenu OJ, Nestadt G, Samuels JF, Costa, PT, Howard WT, Eaton WW. Phobic, panic, and major depressive disorders and the five-factor model of personality. *J Nerv Ment Dis* 2001b; 189 (3): 154–161.
- Bienvenu OJ, Samuels JF, Costa PT, Reti IM, Eaton WW, Nestadt G. Anxiety and depressive disorders and the five-factor model of personality: a higher- and lower-order personality trait investigation in a community sample. *Depress Anxiety* 2004; 20: 92–97.
- Bienvenu OJ, Stein MB, Samuels JF, Onyike CU, Eaton WW, Nestadt G. Personality disorder traits as predictors of subsequent first-onset panic disorder or agoraphobia. *Compr Psychiatry* 2009; 50: 209–214.
- Blatt SJ. Levels of object representation in anaclitic and introjective depression. *Psychoanal Study Child* 1974; 29: 107–157.
- Bornstein RF, Languirand A, Geiselman KJ, Creighton JA, West MA, Gallagher HA, Eisenhart EA. Construct validity of the Relationship Profile Test: A self-report measure of dependency-detachment. *J Pers Assess* 2003; 80: 67–74.
- Bradwejn J, Ahokas A, Stein DJ, Salinas E, Emilien G, Whitaker T. Venlafaxine extended-release capsules in panic disorder: flexible-dose, double-blind, placebo-controlled study. *Br J Psychiatry* 2005; 187: 352–359.
- Brandes M, Bienvenu OJ. Personality and anxiety disorders. *Curr Psychiatry Rep* 2006; 8 (4): 263–269.
- Brown TA, Chorpita BF, Barlow DH. Structural relationships among dimensions of the DSM-IV anxiety and mood disorders and dimensions of negative affect, positive affect, and autonomic arousal. *J Abnorm Psychol* 1998; 107 (2): 179–192.
- Brown GL, Ebert MH, Goyer PF, Jimerson DC, Klein WJ, Bunney WE, Goodwin FK. Aggression, suicide, and serotonin: relationships to CSF amine metabolites. *Am J Psychiatry* 1982; 139 (6): 741–746.
- Brown GL, Goodwin FK, Ballenger JC, Goyer PF, Major LF. Aggression in humans correlates with cerebrospinal fluid amine metabolites. *Psychiatry Res* 1979; 1 (2): 131–139.
- Bruce SE, Yonkers KA, Otto MW, Eisen JL, Weisberg RB, Pagano M, Shea MT, Keller MB. Influence of psychiatric comorbidity on recovery and recurrence in generalized anxiety disorder, social phobia, and panic disorder: a 12-year prospective study. *Am J Psychiatry* 2005; 162: 1179–1187.
- Caci H, Askenazy F, Frequelin N, Nadalet L, Myquel M, Staccini P, Cherick F, Boyer P. Validation of the Impulsivity Rating Scale and relationship with anxiety in healthy French adolescents. *Int J Methods in Psychiatr Res* 1998; 7 (3): 128–135.
- Candilis PJ, McLean RY, Otto MW, Manfro GG, Worthington JJ 3rd, Penava SJ, Marzol PC, Pollack MH. Quality of life in patients with panic disorder. *J Nerv Ment Dis* 1999; 187 (7): 429–434.

- Cannon DM, Klaver JM, Klug SA, Carlson PJ, Luckenbaugh DA, Ichise M, Drevets WC. Gender-specific abnormalities in the serotonin transporter system in panic disorder. *Int J Neuro-psychopharmacol* 2013; 16: 733–743.
- Carrera M, Herrán A, Ramírez ML, Ayestaran A, Sierra-Biddle D, Hoyuela F, Rodríguez – Cabo B, Vázquez-Barquero JL. Personality traits in early phases of panic disorder: implications on the presence of agoraphobia, clinical severity and short-term outcome. *Acta Psychiatr Scand* 2006; 114: 417–125.
- Caspi A, Shiner RL. Personality development. In: Damon W, Lerner R, Eisenberg N, editors. *Handbook of Child Psychology: Social, Emotional, and Personality Development*. 6th ed. New York: Wiley; 2006: 300–365.
- Chavira DA, Stein MB, Golinelli D, Sherbourne CD, Craske MG, Sullivan G, Bystritsky A, Roy-Byrne PP. Predictors of Clinical Improvement in a Randomized Effectiveness Trial for Primary Care Patients with Panic Disorder. *J Nerv Ment Dis* 2009; 197 (10): 715–721.
- Chen YW, Dilsaver SC. Comorbidity of panic disorder in bipolar illness: evidence from the Epidemiologic Catchment Area Survey. *Am J Psychiatry* 1995; 152 (2): 280–282.
- Chopik WJ, Kitayama S. Personality change across the life span: Insights from a cross-cultural, longitudinal study. *J Pers* 2018; 86 (3): 508–521.
- Clark DM. A cognitive approach to panic. *Behav Res Ther* 1986; 24: 461–470.
- Clark LA, Vittengl J, Kraft JD, Jarrett RB. Separate Personality Traits From States to Predict Depression. *J Pers Disord* 2003; 17 (2): 152–172.
- Clark LA, Watson D. Temperament: A new paradigm for trait psychology. In: Pervin LA, John OP, editors. *Handbook of Personality: Theory and research*, 2nd ed. New York: Guilford; 1999: 399–423.
- Clark LA, Watson D, Mineka S. Temperament, personality, and the mood and anxiety disorders. *J Abnorm Psychol* 1994; 103 (1): 103–116.
- Coccaro EF, Siever LJ, Klar HM, Maurer G, Cochrane K, Cooper TB, et al. Serotonergic studies in patients with affective and personality disorders. Correlates with suicidal and impulsive aggressive behavior. *Arch Gen Psychiatry* 1989; 46 (7): 587–599.
- Corchs F, Corregiari F, Ferrão YA, Takakura T, Mathis ME, Lopes AC, Miquel EC, Bernik M. Personality traits and treatment outcome in obsessive-compulsive disorder. *Braz J Psychiatry* 2008; 30 (3): 246–250.
- Costa PT Jr, McCrae RR. *NEO-PI-R. Professional manual* 1992a Odessa, FL: Psychological Assessment Resources.
- Costa PT, McCrae RR. *Revised NEO Personality Inventory (NEO-PI-R) and NEO Five-Factor Inventory (NEO-FFI) Professional Manual*. 1992b Odessa; FL: Psychological Assessment Resources.
- Costa PT Jr, McCraeRR. *The NEO Personality Inventory manual* 1985, Odessa, FL: Psychological Assessment Resources.
- Cougle JR, Timpano KR, Sachs-Ericsson N, Keough ME, Riccardi CJ. Examining the unique relationships between anxiety disorders and childhood physical and sexual abuse in the National Comorbidity Survey-Replication. *Psychiatry Res* 2010; 177 (1–2): 150–155.
- Cuijpers P, van Straten A, Donker M. Personality traits of patients with mood and anxiety disorders. *Psychiatry Res* 2005; 133 (2–3): 229–237.

- Damberg-M, Berggard C, Mattila-Evenden M, et al. Transcription factor AP-2beta genotype associated with anxiety-related personality traits in women. A replication study. *Neuropsychobiology* 2003; 48 (4): 169–175.
- Dammen T, Arensen H, Ekeberg Ø, Friis S. Panic disorder in chest pain patients referred for cardiological outpatient investigation. *J Intern Med* 2001; 245 (5): 497–507.
- de Ruiter C, Cohen L. Personality in panic disorder with agoraphobia: a Rorschach study. *J Pers Assess* 1992; 59 (2): 304–316.
- Del Carlo A, Benvenuti M, Fornaro M, Toni C, Rizzato S, Swann AC, Dell'Osso L, Perugi G. Different measures of impulsivity in patients with anxiety disorders: a case control study. *Psychiatry Res* 2012; 197 (3): 231–236.
- Diagnostic and statistical manual of mental disorders* (3rd Edition) (DSM-III). American Psychiatric Association, Washington DC; 1980
- Diamond DB. Psychotherapeutic approaches to the treatment of panic attacks, hypochondriasis and agoraphobia. *Br J Med Psychol* 1987; 60: 79–84.
- Dickman SJ. Functional and dysfunctional impulsivity: Personality and cognitive correlates. *J Pers Soc Psychol* 1990; 58: 95–102.
- Doerfler LA, Connor DF, Volungis AM, Toscano PF Jr. Panic disorder in clinically referred children and adolescents. *Child Psychiatry Hum Dev* 2007; 3 (1): 57–71.
- Dreessen L, Arntz A. The impact of personality disorders on treatment outcome of anxiety disorders: Best-evidence synthesis. *Behaviour Research and Therapy* 1998; 36: 483–504.
- Dreessen L, Arntz A, Luttels C, Sallaerts S. Personality disorders do not influence the results of cognitive behavior therapies for anxiety disorders. *Comprehensive Psychiatry* 1994; 35 (4): 265–274.
- Du L, Bakish D, Ravindran AV, Hrdina PD. Does fluoxetine influence major depression by modifying five-factor personality traits? *J Affect Disord* 2002; 71 (1–3): 235–241.
- Eaton WW, Kessler RC, Wittchen HU, Magee WJ. Panic and panic disorder in the United States. *Am J Psychiatry* 1994; 151 (3): 413–420.
- Elfhag K. Personality correlates of obese eating behaviour: Swedish universities Scales of Personality and the Three Factor Eating Questionnaire. *Eat Weight Disord* 2005; 10 (4): 210–215.
- Enns MW, Cox BJ. Personality dimensions and depression: review and commentary. *Can J Psychiatry*. 1997; 42 (3): 274–284.
- Eysenck HJ, Eysenck MW. *Personality and individual differences. A natural science approach*. New York: Plenum, 1985.
- Eysenck HJ, Eysenck SBG. *Psychoticism as a dimension of personality*. New York: Crane, Russak, 1976.
- Fagerberg T, Söderman E, Gustavsson P, Agartz I, Jönsson EG. Personality traits in established schizophrenia: aspects of usability and differences between patients and controls using the Swedish universities scales of personality. *Nordic Journal of Psychiatry* 2016; 70 (6): 462–469.
- Fagerberg T, Söderman E, Gustavsson P, Agartz I, Jönsson EG. Stability of personality traits over a five-year period in Swedish patients with schizophrenia spectrum disorder and non-psychotic individuals: a study using the Swedish universities scales of personality. *BMC Psychiatry* 2018; 18 (1): 54.
- Faravelli C, Paterniti S, Scarpato A. 5-year prospective, naturalistic follow-up study of panic disorder. *Compr Psychiatry* 1995; 36 (4): 271–277.

- Fleet RP, Dupuis G, Marchand A, Burelle D, Arsenault A, Beitman BD. Panic disorder in emergency department chest pain patients: prevalence, comorbidity, suicidal ideation, and physician recognition. *Am J Med* 1996; 101: 371–380.
- Foot M, Koszycki D. Gender differences in anxiety-related traits in patients with panic disorder. *Depress Anxiety* 2004; 20 (3): 123–130.
- Fraley RC, Roberts BW. Patterns of continuity: a dynamic model for conceptualizing the stability of individual differences in psychological constructs across the life course. *Psychol Rev* 2005; 112 (1): 60–74.
- Freire RC, Lopes FL, Veras AB, Valença AM, Mezzasalma MA, Nascimento I, Nardi AE. Personality traits spectrum in panic disorder and major depression. *Braz J Psychiatry* 2007; 29 (1): 31–34.
- Freire RC, Zugliani MM, Garcia RF, Nardi AE. Treatment-resistant panic disorder: a systematic review. *Expert Opin Pharmacother* 2016; 17 (2): 159.
- Furukawa TA, Shear MK, Barow DH, Gorman JM, Woods SW, Money R, Etschel E, Engel RR, Leucht S. Evidence-based Guidelines for Interpretation of the Panic Disorder Severity Scale. *Depress Anxiety* 2009; 26: 922–929.
- García LF, Aluja A, García Ó, Cuevas L. Is Openness to Experience an independent personality dimension? Convergent and discriminant validity of the Openness domain and its NEO-PI-R facets. *J Indiv Differ* 2005; 26: 132–138.
- George DT, Hibbeln JR, Ragan PW, Umhau JC, Phillips MJ, Doty L, Hommer D, Rawlings RR. Lactate-induced rage and panic in a select group of subjects who perpetrate acts of domestic violence. *Biol Psychiatry* 2000; 47 (9): 804–812.
- Gerrig RJ, Zimbardo PG. *Psychology and life*. 16th ed. Boston, MA: Allyn and Bacon; 2002.
- Gingnell M, Comasco E, Orelund L, Fredrikson M, Sundström-Poromaa I. Neuroticism-related personality traits are related to symptom severity in patients with premenstrual dysphoric disorder and to the serotonin transporter gene-linked polymorphism 5-HTTLPR. *Arch Women Ment Hlth* 2010; 13 (5): 417–423.
- Goddard AW, Brouette T, Almai A, Jetty P, Woods SW, Charney D. Early coadministration of clonazepam with sertraline for panic disorder. *Arch Gen Psychiatry* 2001; 58: 681–686.
- Goisman R, Warshaw M, Peterson L. Panic, agoraphobia, and panic disorder with agoraphobia. *J Nerv Ment Dis* 1994; 182: 72–79.
- Goodwin RD, Fergusson DM, Horwood LJ. Childhood abuse and familial violence and the risk of panic attacks and panic disorder in young adulthood. *Psychol Med*. 2005; 35: 881–890.
- Goodwin RD, Gotlib IH. Panic attacks and psychopathology among youth. *Acta Psychiatr Scand* 2004; 109: 216–221.
- Gorman JM, Kent JM, Sullivan GM, Coplan JD. Neuroanatomical hypothesis of panic disorder, revised. *Am J Psychiatry* 2000; 157 (4): 493–505.
- Gorman JM, Liebowitz MR, Fyer AJ, Stein J. A neuroanatomical hypothesis for panic disorder. *Am J Psychiatry* 1989; 146 (2): 148–161.
- Gould RA, Otto MW, Pollack MH. A meta-analysis of treatment outcome for panic disorder. *Clin Psychol Rev* 1995; 15: 819–844.
- Grant BF, Hasin DS, Stinson FS, Dawson DA, Patricia Chou S, June Ruan W, Huang B. Co-occurrence of 12-month mood and anxiety disorders and personality disorders in the US: results from the national epidemiologic survey on alcohol and related conditions. *J Psychiatr Res* 2005; 39 (1): 1–9.

- Gustavsson JP. *Validity and stability of self-reported personality traits. Contributions to the evaluation of the Karolinska Scales of Personality (PhD thesis)*. Stockholm: Karolinska Institutet, 1997.
- Gustavsson JP, Bergman H, Edman G, Ekselius L, von Knorring L, Linder J. Swedish universities Scales of Personality (SSP): Construction, internal consistency and normative data. *Acta Psychiatr Scand* 2000; 102 (3): 217–225.
- Gustavsson JP, Linder J, Jönsson E, Weinryb RM. Construction and assessment of five health-relevant personality traits from a Five-Factor Model perspective. *Pers Individ Differ* 2003; 35: 69–89.
- Gustavsson JP, Weinryb RM, Goransson S, Pedersen NL, Asberg M. Stability and predictive ability of personality traits across 9 years. *Pers Individ Differ* 1997; 22: 783–791.
- Guy W. *ECDEU Assessment manual for Psychopharmacology*. Revised. Washington, DC: US Government Printing Office, 1976 DHEW publication ADM 76–338.
- Hamilton M. The assessment of anxiety states by rating. *Br J Med Psychol* 1959; 32: 50–55.
- Hedman E, Andersson G, Lindefors N, Gustavsson P, Lekander M, Ruck C, Andersson E, Ljotsson B. Personality change following internet-based cognitive behavior therapy for severe health anxiety. *PLoS One* 2014; 9 (12): e11387
- Hirschfeld RM, Klerman GL. Personality attributes and affective disorders. *Am J Psychiatry* 1979; 136 (1): 67–70.
- Hirshfeld-Becker DR, Micco J, Henin A, Bloomfield A, Biederman J, Rosenbaum J. Behavioral inhibition. *Depress Anxiety* 2008; 25 (4): 357–367.
- Hirvonen J, Tuominen L, Nägren K, Hietala J. Neuroticism and serotonin 5-HT_{1A} receptors in healthy subjects. *Psychiatry Res* 2015; 234 (1): 1–6.
- Hofmann SG, Shear MK, Barlow DH, Gorman JM, Hersherberger D, Patterson M, Woods SW. Effects of panic disorder treatments on personality disorder characteristics. *Depress Anxiety* 1998; 1: 14–28.
- Iliadis SI, Comasco E, Hellgren C, Kollia N, Sundström Poromaa I, Skalkidou A. Associations between a polymorphism in the hydroxysteroid (11-beta) dehydrogenase 1 gene, neuroticism and postpartum depression. *J Affect Disord* 2017; 207: 141–147.
- Iliadis SI, Koulouris P, Gingnell M, Sylvén SM, Sundström-Poromaa I, Ekselius L, Papadopoulos FC, Skalkidou A. Personality and risk for postpartum depressive symptoms. *Arch Womens Ment Health* 2015; 18 (3): 539–546.
- Izci F, Gültekin BK, Sağlam S, Koc MI, Zincir SB, Atmaca M. Temperament, character traits, and alexithymia in patients with panic disorder. *Neuropsychiatr Dis Treat* 2014; 10: 878–885.
- Itekani T, Kiriike N, Stein MB, Nagao K, Nagata T, Minamikawa N, Shidao A, Fukuhara H. Relationship between perfectionism, personality disorders and agoraphobia in patients with panic disorder. *Acta Psychiatr Scand* 2002; 106: 171–178.
- Johnson JG, Cohen P, Kasen S, Brook JS. Personality disorders evident by early adulthood and risk for anxiety disorders during middle adulthood. *J Anxiety Disord* 2006; 20:408–426.
- Jönsson EG, Ivo R, Gustavsson JP, Geijer T, Forslund K, Mattila-Evenden M. Association between a promoter dopamine D2 receptor gene variant and the personality trait detachment. *Biol Psychiatry* 2003; 53: 577–584.

- Kallasmaa T, Allik J, Realo A, McCrae RR. The Estonian version of the NEO-PI-R: An examination of universal and culture-specific aspects of the five-factor model. *Eur J Pers* 2000; 14: 265–278.
- Kandler C, Bleidorn W, Riemann R, Spinath FM, Thiel W, Angleitner A. Sources of cumulative continuity in personality: a longitudinal multiple-rater twin study. *J Pers Soc Psychol* 2010; 98 (6): 995–1008.
- Katherndahl DA, Realini JP. Where do panic attack sufferers seek care? *J Fam Pract* 1995; 40 (3): 237–243.
- Katerndahl DA, Realini JP. Quality of life and panic-related work disability in subjects with infrequent panic and panic disorder. *J Clin Psychiatry* 1997; 58 (4): 153–158.
- Katon WJ. Clinical practice. Panic disorder. *N Engl J Med* 2006; 354 (22): 2360–2367.
- Katon W, Vitaliano PP, Russo J, Jones M, Anderson K. Panic disorder. Spectrum of severity and somatization. *J Nerv Ment Dis* 1987; 175 (1): 12.
- Katschnig H, Amering M. The long-term course of panic disorder and its predictors. *J Clin Psychopharmacol* 1998; 18 (2): 6S–11S.
- Keller MB, Yonkers KA, Warshaw MG et al. Remission and relapse in subjects with panic disorder and panic with agoraphobia. *J Nerv Ment Dis*. 1994; 182: 290–296.
- Kennedy BL, Schwab JJ, Hyde JA. Defense styles and personality dimensions of research subjects with anxiety and depressive disorders. *Psychiatr Q* 2001; 72: 251–262.
- Kessler RC, Chiu WT, Jin R, Ruscio AM, Shear K, Walters EE. The epidemiology of panic attacks, panic disorder, and agoraphobia in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2006; 63: 415–424.
- Kessler RC, Stang PE, Wittchen HU, Ustun TB, Roy-Burne PP, Walters EE. Lifetime panic-depression comorbidity in the National Comorbidity Survey. *Arch Gen Psychiatry* 1998; 55 (9): 801–808.
- King M, Nazareth I, Levy G, Walker C, Morris R, Weich S et al. Prevalence of common mental disorders in general practice attendees across Europe. *Br J Psychiatry* 2008; 192 (5): 362–367.
- Kipper L, Wachleski C, Salum GA, Heldt E, Blaya C, Manfro GG. Can psychopharmacological treatment change personality traits in patients with panic disorder? *Braz J Psychiatry* 2009; 31 (4): 307–313.
- Klein DF. False suffocation alarms, spontaneous panics, and related conditions. An integrative hypothesis. *Arch Gen Psychiatry* 1993; 50 (4): 306–317.
- Klein DN, Kotov R, Bufferd SJ. Personality and depression: explanatory models and review of the evidence. *Annu Rev Clin Psychol* 2011; 7: 269–295.
- Klerman GL, Weissman MM, Ouellette R, Johnson J, Greenwald S. 1991. Panic attacks in the community. Social morbidity and health care utilization. *JAMA* 265: 742–746.
- Klinterberg B, Schalling D, Edman G, Orelund L, Åsberg M. Personality correlates of platelet monoamine oxidase (MAO) activity in female and male subjects. *Neuropsychobiology* 1987; 18: 89–96.
- Krueger RF, Johnson W. Behavioral genetics and personality: a new look at the integration of nature and nurture. In: John OP, Robins RW, Pervin LA, editors. *Handbook of Personality: Theory and Research*. 3rd ed. New York: Guilford; 2008. 287–310.
- Krueger RF, Tackett JL. Personality and psychopathology: working toward the bigger picture. *J Pers Disord* 2003; 17 (2): 109–128.
- Lahey BB. Public Health Significance of Neuroticism. *Am Psychol* 2009; 64 (4): 241–256.

- Lamers F, van Oppen P, Comijs HC, Smit JH, Spinhoven P, van Balkom AJ et al. Comorbidity patterns of anxiety and depressive disorders in a large cohort study: the Netherlands Study of Depression and Anxiety (NESDA). *J Clin Psychiatry* 2011; 72 (3): 341–348.
- Lecrubier Y, Braconnier A, Said S, Payan C. The impulsivity rating scale (IRS): preliminary results. *Eur Psychiatry* 1995; 10: 331–338.
- Lee JH, Dunner DL. The effect of anxiety disorder comorbidity on treatment resistant bipolar disorders. *Depress Anxiety* 2008; 25 (2): 91–97.
- Lee EH, Kim JH, Yu BH. Reliability and validity of the self-report version of the Panic Disorder Severity Scale in Korea. *Depress Anxiety* 2009; 26 (8) E120–123. doi: 10.1002/da.20461.
- Liebowitz MR. Panic disorder as a chronic illness. *J Clin Psychiatry* 1997; 58 (13): 5–8.
- Lydiard RB. Assessment and Management of Treatment-Resistance in Panic Disorder. *FOCUS* 2011; 9 (3): 253–263.
- McCrae RR, Costa PT. Personality trait structure as a human universal. *Am Psychol* 1997; 52: 509–516.
- McNally RJ. Anxiety sensitivity and panic disorder. *Biol Psychiatry* 2002; 52 (10): 938–946.
- Marchesi C, Cantoni A, Fontò S, Giannelli MR, Maggini C. The effect of pharmacotherapy on personality disorders in panic disorder: a one year naturalistic study. *J Affect Disord* 2005; 89 (1–3): 189–194.
- Marchesi C, Cantoni A, Fontò S, Giannelli MR, Maggini C. The effect of temperament and character on response to selective serotonin reuptake inhibitors in panic disorder. *Acta Psychiatr Scand* 2006b; 114 (3): 203–210.
- Marchesi C, De Panfilis C, Cantoni A, Fontò S, Giannelli MR, Maggini C. Personality disorders and response to medication treatment in panic disorder: a 1-year naturalistic study. *Prog Neuropsychopharmacol Biol Psychiatry* 2006a; 30 (7): 1240–1245.
- Marchesi C, De Panfilis C, Cantoni A, Giannelli MR, Maggini C. Effect of pharmacological treatment on temperament and character in panic disorder. *Psychiatry Res* 2008; 158 (2): 147–154.p-v
- Marchevsky D. Selective serotonin reuptake inhibitors and personality change. *Br J Psychiatry* 1999; 175: 589–590.
- Maron E, Hettema JM, Shlik J. Advances in molecular genetics of panic disorder. *Mol Psychiatry* 2010a; 15 (7): 681–701.
- Maron E, Tõru I, Hirvonen J, Tuominen L, Lumme V, Vasar V, Shlik J, Nutt DJ, Helin S, Nagren K, Tiitonen J, Hietala J. Brain serotonin transporter availability in panic disorder: gender differences and implications for treatment response to escitalopram. *J Psychopharmacol* 2010b; 25: 952–959.
- Martin R, Watson D, Wan CK. A three-factor model of trait anger: Dimensions of affect, behavior, and cognition. *J Pers* 2000; 68: 869–897.
- Masi G, Favilla L, Mucci M, Millepiedi S. Panic disorder in clinically referred children and adolescents. *Child Psychiat Hum Devel* 2000; 31: 139–151.
- McCrae RR, Costa PT Jr. Personality trait structure as a human universal. *American Psychologist* 1997; 52, 509–516.
- Mendlowicz MV, Stein MB. Quality of life in individuals with anxiety disorders. *Am J Psychiatry* 2000; 157 (5): 669–682.

- Mennin DS, Heimberg RG. The impact of comorbid mood and personality disorders in the cognitive-behavioral treatment of panic disorder. *Clin Psychol Rev* 2000; 20 (3): 339–357.
- Mitte K. A meta-analysis of the efficacy of psycho- and pharmacotherapy in panic disorder with and without agoraphobia. *J Affect Disord* 2005; 88: 27–45.
- Morey LC, Shea MT, Markowitz JC, Stout RL, Hopwood CJ, Gunderson JG, Grilo CM, McGlashan TH, Yen S, Sanislow CA, Skodol AE. State effects of major depression on the assessment of personality and personality disorder. *Am J Psychiatr* 2010; 167 (5): 528–535.
- Moscovitch DA, McCabe RE, Antony MM, Rocca L, Swinson RP. Anger experience and expression across the anxiety disorders. *Depress Anxiety* 2008; 25 (2): 107–113.
- Muller J, Handlin L, Harlén M, Lindmark U, Ekström A. Mechanical massage and mental training programmes affect employees' anxiety, stress susceptibility and detachment – a randomised explorative pilot study. *BMC Complement Altern Med* 2015; 15: 302.
- Na HR, Kang EH, Lee JH, Yu BH. The Genetic Basis of Panic Disorder. *J Korean Med Sci* 2011; 26 (6): 701–710.
- Nash JR, Sargent PA, Rabiner EA, Hood SD, Argyropoulos SV, Potokar JP, et al. Serotonin 5-HT_{1A} receptor binding in people with panic disorder: positron emission tomography study. *Br J Psychiatry* 2008; 193 (3): 229–234.
- Navarro B, Sánchez M, Herrán A, Sierra-Biddle D. Relationship between personality traits and panic disorder. *Actas Esp Psiquiatr* 2013; 41 (1): 27–32.
- Neuger J, Wistedt B, Aberg-Wistedt A, Stain-Malmgren R. Effect of citalopram treatment on relationship between platelet serotonin functions and the Karolinska scales of personality in panic patients. *J Clin Psychopharmacol* 2002; 22 (4): 400–405.
- Niazi S, Adil A. Relationship between panic attacks and aggression with respect to age and gender. *J Indian Acad Appl Psychol* 2008; 34 (2): 287–293.
- Noyes R Jr, Crowe RR, Harris EL, Hamra BJ, McChesney CM, Chaudhry DR. Relationship between panic disorder and agoraphobia. A Family Study. *Arch Gen Psychiatry* 1986; 43: 227–232.
- Noyes R, Reich J, Christiansen J, Suelzer M, Pfohl B, Coryell WA. 1990. Outcome of panic disorder: Relationship to diagnostic subtypes and comorbidity. *Arch Gen Psychiatry* 47: 809–818.
- O'Keane VO, Moloney E, O'Neill H, O'Connor A, Smith C, Dinan TG. Blunted prolactin responses to d-fenfluramine in sociopathy. Evidence for subsensitivity of central serotonergic function. *Br J Psychiatry* 1992; 160: 643–646.
- Ormel J, Jeronimus B, Kotov R, Riese H, Bos E, Hankin B. Neuroticism and common mental disorders: Meaning and utility of a complex relationship. *Clin Psychol Rev* 2013; 33 (5): 686–697.
- Ortet G, Ibáñez MI, Llerena A, Torrubia R. Underlying traits of the Karolinska Scales of Personality (KSP). *Eur J Psychol Assess* 2002; 18: 139–148.
- Otto MW, Tuby KS, Gould RA, McLean RY, Pollack MH. An effect-size analysis of the relative efficacy and tolerability of serotonin selective reuptake inhibitors for panic disorder. *Am J Psychiatry* 2001; 158: 1989–1992.
- Ozkan M, Altindag A. Comorbid personality disorders in subjects with panic disorder: do personality disorders increase clinical severity? *Compr Psychiatry* 2005; 46 (1): 20–26.

- Park HU, Kim EJ, Ku JI, Woo JM, Lee SH, Chung SK, Lee SY, Kim YR, Kang EH, LeeJH, Yu BH. Psychological characteristics of early remitters in patients with panic disorder. *Psychiatry Res* 2012; 197 (3): 237–241.
- Pelissolo A, Lepine JP. Epidemiology of depression and anxiety disorders. In: Montgomery SA, den Boer JA, editors. *SSRI's in depression and anxiety*. New York: Wiley & Sons Inc. 1998.
- Perna G. Panic disorder: from psychopathology to treatment. *J Psychopathol* 2012; 18: 82–91.
- Pollack MH, Otto MW. Long-term course and outcome of panic disorder. *J Clin Psychiatry* 1997; 58 (2): 57–60.
- Pollack MH, Otto MW, Worthington JJ, Manfro GG, Wolkow R. Sertraline in the treatment of panic disorder: a flexible-dose multicenter trial. *Arch Gen Psychiatry* 1998; 55 (11): 1010–1016.
- Powers A, Western D. Personality subtypes in patients with panic disorder. *Compr Psychiatry* 2009; 50 (2): 164–172.
- Practice guideline for the treatment of patients with panic disorder. Work Group on Panic Disorder. American Psychiatric Association. *Am J Psychiatry* 1998; 155 (5 Suppl): 1–34.
- Prasko J, Houbová P, Novák T, Záleský R, Espa-Cervená K, Pasková B, Vyskocilová J. Influence of personality disorder on the treatment of panic disorder – comparison study. *Neuro Endocrinol Lett* 2005; 26 (6): 667–674.
- Preti A, Vrublevska J, Veroniki AA, Huedo-Medina TB, Kyriazis O, Fountoulakis KN. Prevalence and treatment of panic disorder in bipolar disorder: systematic review and meta-analysis. *Evid Based Ment Health* 2018; 21 (2): 53–60.
- Pull CB, Damsa C. Pharmacotherapy of panic disorder. *Neuropsychiatr Dis Treat* 2008; 4 (4): 779–795.
- Radu D, Ahlin A, Svanborg P, Lindefors N. Pentagastrin test for anxiety – psychophysiology and personality. *Psychopharmacology (Berl)* 2003; 166 (2): 139–145.
- Rasul A, El-Nour H, Lonne-Rahm SB, Fransson O, Johansson C, Johansson B, Zubeidi M, Seeberg E, Djurfeldt DR, Azmitia AE, Nordlind K. Serotonergic Markers in Atopic Dermatitis. *Acta Derm Venereol* 2016; 96 (6): 732–736.
- R Core Team (2014). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Regier DA, Rae DS, Narrow WE, Kaelber CT, Schatzberg AF. Prevalence of anxiety disorders and their comorbidity with mood and addictive disorders. *Br J Psychiatry* Suppl. 1998; 173: 24–28.
- Reich JH, Green AI. Effect of personality disorders on outcome of treatment. *J Nerv Ment Dis* 1991; 179 (2): 74–82.
- Reich J, Noyes R Jr, Hirshfeld R, Coryell W, O’Gorman T. State and personality in depressed and panic patients. *Am J Psychiatry* 1987; 144 (2): 181–187.
- Reznick JS, Hegeman IM, Kaufman ER, Woods SW, Jacobs M. Retrospective and concurrent self-report of behavioral inhibition and their relation to adult mental health. *Development and Psychopathology* 1992; 4 (2): 301–321.
- Rief W, Martin A, Klaiberg A, et al. Specific effects of depression, panic, and somatic symptoms on illness behavior. *Psychosom Med* 2005; 67: 596–601.
- Remröd C, Sjöström K, Svensson Å. Subjective stress reactivity in psoriasis – a cross sectional study of associated psychological traits. *BMC Dermatol* 2015; 15: 6.

- Roberts BW, Luo J, Briley DA, Chow PI, Su R, Hill PL. A systematic review of personality trait change through intervention. *Psychol Bull* 2017; 143 (2): 117–141.
- Robins LN, Reiger DA.(eds.) *Psychiatric disorders in America. The Epidemiologic Catchment Area Study*. 1991; New York: The Free Press.
- Rocca WA, Grossardt BR, Peterson BJ, Bower JH, Trenerry MR, Ahlskog JE, Sanft KR, de Andrade M, Maraganore DM. The Mayo Clinic cohort study of personality and aging: design and sampling, reliability and validity of instruments, and baseline description. *Neuroepidemiology* 2006; 26 (3): 119–129.
- Rothbart MK, Bates JE. Temperament. In Damon W, Lerner R, Eisenberg N (Eds.), *Handbook of child psychology: Vol. 3. Social, emotional, and personality development*. 6th ed., 2006; New York: Wiley: 99–166.
- Roy-Byrne PP, Craske MG, Stein MB. Panic disorder. *Lancet* 2006; 368: 1023–1032.
- Roy-Byrne PP, Geraci M, Uhde TW. Life events and the onset of panic disorder. *Am J Psychiatry* 1986; 143 (11): 1424–1427.
- Santor DA, Bagby RM, Joffe RT. Evaluating stability and change in personality and depression. *J Pers Soc Psychol* 1997; 73: 1354–1362.
- Schalling D. Psychopathy-related personality variables and the psychophysiology of socialization. In: *Psychopathic behaviour. Approaches to research*, Hare RD, Schalling D (Eds.), 1978 Chichester: John Wiley and Sons; 85–106.
- Schalling D, Åsberg M, Edman G, Orelund L. Markers for vulnerability to psychopathology: Temperament traits associated with platelet MAO activity. *Acta Psychiatr Scand* 1987; 76: 172–182.
- Schmidt NB, Lerew DR, Trakowski JH. Body vigilance in panic disorder: evaluating attention to bodily perturbations. *J Consult Clin Psychol* 1997; 65 (2): 214–220.
- Scocco P, Barbieri I, Frank E. Interpersonal problem areas and onset of panic disorder. *Psychopathology* 2007; 40: 8–13.
- Seivewright H, Tyrer P, Johnson T. Change in personality status in neurotic disorders. *Lancet* 2002; 359 (9352): 2253–2254.
- Seki Y, Nagata S, Shibuya T, Yoshinaga N, Yokoo M, Ibuki H, Minamitani N, Kusunoki M, Inada Y, Kawasoe N, Adachi S, Yoshimura K, Nakazato M, Iyo M, Nakagawa A, Shimizu E. A feasibility study of the clinical effectiveness and cost-effectiveness of individual cognitive behavioral therapy for panic disorder in a Japanese clinical setting: an uncontrolled pilot study. *BMC Res Notes* 2016; 9: 458.
- Shtasel DL, Gur RE, Mozley PD, Richards J, Taleff MM, Heimberg C, Gallacher F, Gur RC. Volunteers for biomedical research. Recruitment and screening of normal controls. *Arch Gen Psychiatry* 1991; 48: 1022–1025.
- Shear MK, Brown TA, Barlow DH, Money R, Sholomskas DE, Woods SW, Gorman JM, Papp LA. Multicenter collaborative panic disorder severity scale. *Am J Psychiatry* 1997; 154 (11): 1571–1575.
- Shear MK, Rucci P, Williams J, Frank E, Grochocinski V, Vander Bilt J, Houck P, Wang T. Reliability and validity of the Panic Disorder Severity Scale: replication and extension. *J Psychiatr Res* 2001; 35: 293–296.
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R, Dunbar GC. The Mini International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview. *J Clin Psychiatry* 1998; 59 (20): 22–33; 34–57.
- Shlik J, Aluoja A, Kihl E. *MINI 5.0.0. Mini rahvusvaheline neuropsühhiaatriiline intervjuu DSM-IV. Estonian version of MINI international neuropsychiatric interview*; 1999.

- Siegel A, Bhatt S, Bhatt R, Zalcman SS. The neurobiological bases for development of pharmacological treatments of aggressive disorders. *Curr Neuropsychopharmacol* 2007; 5 (2): 135–147.
- Silva H, Iturra P, Solari A, Villarroel J, Jerez S, Jimenez M, Galleguillos F, Bustamante ML. Fluoxetine response in impulsive-aggressive behavior and serotonin transporter polymorphism in personality disorder. *Psychiatr Genet* 2010; 20 (1): 25–30.
- Simpson RJ, Kazmierczak T, Power KG, Sharp DM. Controlled comparison of the characteristics of patients with panic disorder. *Br J Gen Pract* 1994; 44: 352–356.
- Skodol AE., Oldham JM, Hyler SE, Stein DJ, Hollander E, Gallaher PE, Lopez AE. Patterns of anxiety and personality disorder comorbidity. *J Psychiatr Res* 1995; 29: 361–374.
- Slaap BR, den Boer JA. The prediction of nonresponse to pharmacotherapy in panic disorder: a review. *Depress Anxiety* 2001; 14: 112–122.
- Stalenheim EG. Long-term validity of biological markers of psychopathy and criminal recidivism: Follow-up 6-8 years after forensic psychiatric investigation. *Psychiatry Res* 2004; 121: 281–291.
- Sparding T, Pålsson E, Joas E, Hansen S, Landén M. Personality traits in bipolar disorder and influence on outcome. *BMC Psychiatry* 2017; 17: 159.
- Summerfeldt L, Hood K, Antony MM, Richter MA, Swinson RP: Impulsivity in obsessive-compulsive disorder: comparisons with other anxiety disorders and within tic-related subgroups. *Pers Individ Dif* 2004; 36: 539–553.
- Susman J, Klee B. The role of high-potency benzodiazepines in the treatment of panic disorder. *Prim Care Companion J Clin Psychiatry* 2005; 7: 5–11.
- Zanarini MC, Frankenburg FR, Dubo ED, Sickel AE, Trikha A, Levin A, Reynolds V. Axis I comorbidity of borderline personality disorder. *Am J Psychiatry* 1998; 155: 1733–1739.
- Zane RD, McAfee AT, Sherburne S, Billeter G, Barsky A. Panic disorder and emergency services utilization. *Acad Emerg Med* 2003; 10 (10): 1065–1069.
- Zaubler TS, Katon W. Panic disorder in the general medical setting. *J Psychosom Res* 1998; 44 (1): 25–42.
- Zuckerman M, Kuhlman DM, Joireman J, Teta P, Kraft MA. Comparison of three structural models for personality: The Big Three, the Big Five, and the Alternative Five. *J Pers Soc Psychol* 1993; 65: 757–768.
- Zuckerman M, Kuhlman DM, Thornquist M, Kiers H. Five (or three) robust questionnaire scale factors of personality without culture. *Pers Individ Differ* 1991; 12: 929–941.
- Zugliani MM, Martin-Santos R, Nardi AE, Freire RC. Personality traits in panic disorder patients with and without comorbidities. *J Nerv Ment Dis* 2017; 205 (11): 855–858.
- Zun LS. Panic disorder. Diagnosis and treatment in emergency medicine. *Ann Emerg Med* 1997; 30: 92–96.
- Takano A, Arakawa R, Hayashi M, Takahashi H, Ito H, Suhara T. Relationship between neuroticism personality trait and serotonin transporter binding. *Biol Psychiatry* 2007; 62 (6): 588–592.
- Tang TZ, DeRubeis RJ, Hollon SD, Amsterdam J, Shelton R, Schalet B. Personality change during depression treatment: a placebo-controlled trial. *Arch Gen Psychiatry* 2009; 66 (12): 1322–1330.

- Tueth MJ. Managing recurrent nonischemic chest pain in the emergency department. *Am J Emerg Med* 1997; 15: 170–172.
- Tweed JL, Schoenbach VJ, George LK, Blazer DG. The effects of childhood parental death and divorce on six-month history of anxiety disorders. *Br J Psychiatry* 1989; 154: 823–828.
- Tõru I, Aluoja A, Võhma Ü, Raag M, Vasar V, Maron E, Shlik J. Associations between personality traits and CCK-4-induced panic attacks in healthy volunteers. *Psychiatry Res* 2010; 178 (2): 342–347.
- van Apeldoorn FJ, van Hout WJ, Mersch PP, Huisman M, Slaap BR, Hale WW 3rd, Visser S, van Dyck R, den Boer JA. Is a combined therapy more effective than either CBT or SSRI alone? Results of a multicenter trial on panic disorder with or without agoraphobia. *Acta Psychiatr Scand* 2008; 117 (4): 260–270.
- Vanderkooy JD, Kennedy SH, Bagby RM. Antidepressant side effects in depression patients treated in a naturalistic setting: a study of bupropion, moclobemide, paroxetine, sertraline, and venlafaxine. *Can J Psychiatry* 2002; 47 (2): 174–80.
- Volgsten H, Ekselius L, Poromaa IS, Svanberg AS. Personality traits associated with depressive and anxiety disorders in infertile women and men undergoing in vitro fertilization treatment. *Acta Obstet Gynecol Scand* 2010; 89 (1): 27–34.
- Võhma Ü, Aluoja A, Vasar V, Shlik J, Maron E. Evaluation of personality traits in panic disorder using Swedish universities Scales of Personality. *J Anxiety Disord* 2010; 24: 141–146.
- Wachleski C, Salum GA, Blaya C, Kipper L, Paludo A, Salgado AP and Manfro GG. Harm avoidance and self-directedness as essential features of panic disorder patients. *Comp Psychiatry* 2008; 49 (5): 476–481.
- Walderhaug E, Herman AI, Magnusson A, Morgan MJ, Landrø NI. The short (S) allele of the serotonin transporter polymorphism and acute tryptophan depletion both increase impulsivity in men. *Neurosci Lett* 2010; 473 (3): 208–211.
- Weissman MM, Bland RC, Canino GJ, Faravelli C, Greenwald S, Hwu HG, Joyce PR, Karam EG, Lee CK, Lellouch J, Lépine JP, Newman SC, Oakley-Browne MA, Rubio-Stipec M, Wells JE, Wickramaratne PJ, Wittchen HU, Yeh EK. The cross-national epidemiology of panic disorder. *Arch Gen Psychiatry* 1997; 54 (4): 305–309.
- Whiteside SP, Lynam DR. The five factor model and impulsivity: Using a structural model of personality to understand impulsivity. *Pers Individ Differ* 2001; 30: 669–689.
- WHO (ed) *The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines*. Geneva, 1992.
- Wiborg IM, Falkum E, Dahl AA. Is harm avoidance an essential feature of patients with panic disorder? *Compr Psychiatry* 2005; 46: 311–314.
- Wittchen HU, Essau CA. Epidemiology of panic disorder: progress and unresolved issues. *J Psychiatr Res* 1993; 27 (1): 47–68.
- Wittchen HU, Nelson CB, Lachner G. Prevalence of mental disorders and psychosocial impairments in adolescents and young adults. *Psychol Med* 1998; 2: 109–126.
- Working Group on Panic Disorder. Practice guideline for the treatment of patients with panic disorder. *Am J Psychiatry* 1998; 155: Suppl: 1–34.
- Yates WR. Phenomenology and epidemiology of panic disorder. *Ann Clin Psychiatry* 2009; 21: 95–102.

SUMMARY IN ESTONIAN

Paanikahäirega patsientide isikuomaduste, kliiniliste näitajate seos farmakoloogilise ravivastusega

Sissejuhatus

Paanikahäire (PH) kui eraldiseisev häire lisati ametlikult psühhiaatriliste häirete nimistusse 1980. aastal DSM III publitseerimisel (APA, 1980). Paanikahäiret iseloomustavad korduvad paanikahood ning ootusärevus nende hoogude ees, sageli kujuneb sellest välja agorafobia (DSM-IV, RHK-10).

Paanikahäire eluaegne esinemissagedus on 1.5–3.5%, kusjuures 12 kuu esinemissagedus on naistel kõrgem kui meestel (Klerman jt. 1991). PH algab tavaliselt kahekümnendate eluaastate keskel, naistel on leitud kõrgeim haaratus 25–34 eluaasta vahel, meestel 30–44 eluaasta vahel (Wittchen ja Essau, 1993).

PH kahjustab tõsiselt sotsiaalset toimimist, töö ja isikliku eluga toimetulekut, mõjutades seega oluliselt üldist elukvaliteeti (Candilis jt. 1999). PH patsientidel on kalduvus tõlgendada kehalisi sümptomeid katastroofilisel moel, seetõttu on selle häirega isikute meditsiiniteenuste kasutamine ebaproportsionaalselt kõrge (Barsky jt. 1999), võrreldes kontrollvalimiga teevad nad oluliselt sagedamini visiite perearsti vastuvõtule (Simpson jt. 1994), kasutavad sagedamini vältimatut abi ja vaimse tervise abi võrreldes teiste üldmeditsiiniliste ambulatoorsete patsientidega (Barsky jt. 1999). PH loetakse üldiselt kroonilise kuluga häireks. Pikaajases uuringus on kolme aasta järel leitud ainult 10% patsiente täielikult sümptomivabana (Noyes jt. 1990), täielik remissioon viie aasta pärast on leitud vaid 12% PH patsientidest (Faravelli jt. 1995).

Paanikahäire etiopatogeneesis on kompleksne, hõlmates sotsiodemograafilisi, geneetilisi, bioloogilisi ja psühholoogilisi faktoreid (Andrizano 2012, Wittchen ja Essau 1993, Scocco jt. 2007, Faravelli 1995, Tweed jt. 1989; Cogle jt. 2010, Goodwin jt. 2005b, Roy-Byrne jt. 1986; Assellmann jt. 2017, Hirshfeld-Becker jt. 2008). Paanikahäire ravis on kaksikpimedates uuringutes tõestanud efektiivsust viie ravimiklassi preparaadid: selektiivsed serotoniini tagasihaarde inhibiitorid (SSTI), serotoniini noradrenaliini tagasihaarde inhibiitorid, tritsüklilised antidepressandid, monoamiinioksüdaasi inhibiitorid ning bensodiasepiinid (Mitte, 2005; Bradwejn, 2005; Otto jt. 2001; Gould jt. 1995; Susman & Klee 2005; Goddard jt. 2001). Avaldatud uuringu kohaselt 20–40% paanikahäirega patsientidest ei anna ravivastust farmakoteraapiale ja 30–40% kognitiiv-käitumuslikule psühhoteraapiale (van Apeldoorn jt. 2008), seega mitte kõik patsiendid ei saavuta remissiooni.

SSTI-d leiti Otto (2001) ja kolleegide poolt läbi viidud 12 uuringut hõlmavas metaanalüüsis üldistunud ärevushäire ning paanikahoogude sageduse vähendamisel platseebost oluliselt efektiivsemad. Ka käesolevas uuringus on kasutatud paanikahäire raviks SSTI estsitalopraami 10–20 mg päevas.

Isiksust on kirjeldatud kui isiku unikaalset psühholoogiliste omaduse kogumit, mis väljendub läbi elu erinevates olukordades kindlates käitumismustrites (Gerrig

& Zimbardo, 2002). Isiksuse põhijooni on peetud elu jooksul stabiilseks, kuid viimastel aastatel on leitud, et need võivad läbi elu siiski muutuda seoses isiksuse küpsemisega või elusündmuste mõjul (Seviewright jt. 2002, Cohrs jt. 2008). Isiksuse omaduste, eriti neurootilisuse ja ekstravertsuse muutusi, on seostatud ka kliinilise sekkumisega (Roberts jt. 2017). On oletatud, et isiksus mängib otsustavat rolli paanikahäire alguse ning arengu juures (Clark, Watson, & Ineka, 1994). Paanikahäirega patsientidel on leitud kõrgeenenud neurootilisust ja madalat positiivset emotsionaalsust (Bienvenu jt. 2001b; Bienvenu jt. 2004; Carrera jt. 2006; Zugliani jt. 2017). Madalat ekstravertsust on seostatud agorafobiaga, mitte otseselt paanikahäirega (Bienvenu jt. 2004). Samas on leitud, et kõrgeenenud neurootilisus ja madal ekstravertsus tulevad esile pigem komorbiidsuse korral (Cuijpers jt. 2005; Zugliani jt. 2013).

Paanikahäire ja isiksuse joonte seoste osas on mitu autorit avaldanud arvamust, et isiksuse jooni, eriti neurootilisust, võivad mõjutada nii häire ägedad sümptomid, farmakoteraapia kui ka kognitiiv-käitumuslik teraapia (Rocca jt. 2006, Corchs jt. 2008). Võrreldes depressiooniga on ärevushäirete, sealhulgas paanikahäire seost isiksuse omadustega vähem uuritud ja mõistetud (Bienvenu & Brandes, 2005). Neid seoseid on käsitletud kahepoolsetena, kus isiksus võib mõjutada ravitulemust ning ravi võib viia muutustele isiksuse mustriks. On leitud, et isiksuse kõrvalekalded mõjutavad paanikahäire sümptomaatika intensiivsust (Sokol jt. 1995, Ozkan jt. 2005) ja võivad negatiivselt mõjutada ravitulemust (Slaap&den Boer 2001; Marchesi jt. 2006). Samas on uuringuid, kus ükski NEO-PI-R-ga hinnatud isiksuse joon ei olnud seotud häire kliinilise raskuse ega lühiaegse ravitulemusega SSTI-ga (Carrera jt. 2006). Samuti ei leitud seost isiksuse joonte ja PH sümptomite remissiooni vahel (Marchesi jt.2006). Seega on uuringute tulemused isiksuse joonte mõjust paanikahäire ravitulemusele jäänud vastuoluliseks.

Hinnates PH farmakoloogilise ravi mõju isiksusele leiti, et sümptomite paranemisel vähenesid oluliselt ka neurootilisusega seotud isiksuse jooned (Reich jt. 1991; Noyes jt. 1986, Tang jt 2009, Barlow jt. 2013).

Kognitiiv-käitumisteraapia tulemusena leiti, et SSP-ga hinnatud neurootilised jooned vähenesid, samas ekstravertus ja agressiivsus jäid muutumatuks (Hedman jt. 2014). Märkimisväärne on, et kuigi on uuringuid, kus PH remissioon on seotud isiksuse joonte osalise "normaliseerumisega" (Brandes jt. 2006), on mitmeid uuringuid, kus isiksuse jooned jäävad stabiilseks kogu PH kulus (Morey jt. 2010, Santor jt.1997).

Uuringu põhjendus

Paanikahäire ja isiksuse seosed ei ole endiselt lõplikult selgitatud, see on jäänud teadusliku arutelu ja uurimise teemaks, sealjuures uuringute valimid on olnud suhteliselt väikesed ning tulemused vasturääkivad. Ärevushäirete, sealhulgas paanikahäire ravitulemust ennustavate tegurite parem tundmine aitab kaasa personaalmeditsiinile, patsiendile individuaalselt parima ravi valikuvõimalusele.

Isiksusel näib olevat ülioluline roll psüühikahäirete patogeneesis ja prognoosis. Oluline on ka see, et isiksuse omadused ravitulemuse ennustajana on kergesti hinnatavad. Käesolevas töös on kasutatud isiksuse hindamiseks Rootsi ülikoolide isiksuseskaalat (SSP), mille 13 alaskaalat on näidanud häid psühhomeetrilisi omadusi Rootsi normpopulatsioonis, kuid mis on eelkõige disainitud mõõtma isiksuse jooni psühhopatoloogia esinemisel (Schalling 1978; Gustavsson jt. 2000). SSP on tõestanud kasulikkust mitmetes psühhobioloogilistes uuringutes (Damberg jt. 2003, Jönsson jt. 2003). Faktoranalüüsi alusel mõõdab SSP kolme põhifaktorit: neurootilisus, ekstravertsus, agressiivsus (Gustavsson jt. 2000). Käesolevas uuringus kontrolliti SSP eestikeelse versiooni reliaablust ning valiidsust. See võimaldas hinnata SSP faktorite universaalsust originaalpopulatsioonist erinevas kultuuritaustas, samuti SSP suhet viiefaktorilise isiksusemudeliga.

Uurimistöö eesmärgid

Käesoleva töö üldine eesmärk oli kirjeldada paanikahäirega patsientide isiksuse omadusi ja uurida nende isiksuse joonte seoseid ravitulemusega, kasutades valideeritud SSP hindamisskaalat.

Kitsamad eesmärgid:

1. Kontrollida eestikeelse SSP reliaablust ja valiidsust (I)
2. Iseloomustada SSP-ga mõõdetud isiksuse joonte positsiooni viiefaktorilise isiksuse- mudeli suhtes (I)
3. Tuvastada paanikahäirega patsientide ja tervete isikute isiksuse joonte erinevused hinnatuna SSP-ga (II)
4. Uurida isiksuse omaduste erinevusi paanikahäirega patsientidel ilma ja koos afektiivse häirega ning jälgida paanikahäirel SSP domeenide (alaskaalad, üldfaktorid) ning erinevate demograafiliste ja kliiniliste muutujate vahelisi seoseid (II)
5. Uurida estsitalopraami ravi mõju paanikahäirega patsientide isiksuse joontele (III)
6. Leida, kas SSP-ga mõõdetud isiksuse omadused ennustavad paanikahäire farmakoloogilise ravi tulemust

Uuringutes osalejad ja meetodid

I uuringus osales 529 uuritavat. Esimeses grupis oli 331 tervet vabatahtlikku erinevatest uuringuprojektidest Tartu Ülikooli Psühhiaatrikliinikus ning teine grupp koosnes 198 erineva haridusliku ja sotsiaalse taustaga isikust. Mõlemas grupis osalejad täitsid Rootsi ülikoolide isiksuseskaala (SSP) (Gustavsson jt. 2000) ja 197 isikut teisest grupist täitsid NEO-PI-R (Costa &McRae, 1992b, Kallasmaa jt. 2000) eestikeelse variandi.

II uuringus olid kaasatud 193 paanikahäirega patsienti Tartu Ülikooli Psühhiaatrikliinikust ning 314 ajalehekuulutuse kaudu leitud tervet vabatahtlikku.

Paanikahäirega patsientidest 71 olid ainult paanikahäirega, 122 komorbiidse paanikahäirega, kusjuures 93-l esines kaasuvana depressioon, 29-l bipolaarne häire, 46-l teised ärevushäired, sealhulgas obsessiivne-kompulsiivne häire ja sotsiaalfobia. Agorafoobia esines kaasuvalt 115-l, sealhulgas ainult paanikahäirega grupis 46-l ja komorbiidse häirega grupis 69-l. Kõik patsiendid ning tervete kontrollgrupp täitis SSP eestikeelse variandi.

III uuringus osales lõplikus valimis 107 paanikahäirega patsienti, kes täitsid SSP algvisiidil ning 12-nädalase ravi järel 10–20 mg estsitalopraamiga. Kaasuvalt täitsid uuritavad algvisiidil ning iga järgneva kahe nädala järel lisaskaalad PDSS (Shear jt. 1997), CGI (Guy 1976), HAM-A (Hamilton 1959), emotsionaalse enesetunde küsimustiku (EEK-2) (Aluoja jt. 1999), TSES (Vanderkooy jt. 2002).

Kõikides uuringutes tõendati paanikahäire ja teiste häirete olemasolu või puudumist kliinilise intervjuu ning rahvusvahelise neuropsühhiaatrilise intervjuu MINI (M.I.N.I. 5.0.0; Sheehan jt. 1998) läbiviimisega.

Peamised tulemused

Esimeses uuringus olid SSP alaskaaladest naistel oluliselt kõrgemad väärtused psüühilisel ärevusel, vastuvõtlikkusel stressile, kehtestavuse puudumisel ja sotsiaalsel soovitavusel. Meestel olid kõrgemad näitajad verbaalsel ja füüsilisel agressioonil, umbusklikkusel ja seikluste otsimisel. Suurimad soolised erinevused ilmnesisid agressiooni ja ärevuse alaskaaladel.

Rootsi ülikoolide isiksuseskaalal (SSP) leiti kolmefaktoriline struktuur.

Esimene faktor hõlmas skaalasid, mis iseloomustavad neurootilisust, kirjeldades ära 27,8% andmete koguhajuvusest. Kõige suurema faktorlaadungiga selles faktoris olid somaatiline ja füüsiline ärevus, vastuvõtlikkus stressile, kehtestavuse puudumine.

Teine faktor kirjeldas 20,2% andmete koguhajuvusest. Faktori moodustasid alaskaalad verbaalne ja füüsiline agressiivsus, ärrituvus, isoleerumine ja madal sotsiaalne soovitavus, osutades agressioonile ja mittekonformsusele. Kusjuures isoleeritus ja ärritatus haakusid ka esimese ja kolmanda faktoriga.

Kolmas faktor kirjeldas ära 14,8% andmete koguhajuvusest. Kolmandasse faktorisse kuulusid impulsiivsus ja seikluste otsimine.

Umbusklikkus haakus lähedaselt kahe faktoriga, mis mõõtsid neurootilisust ning agressiooni.

Seega faktoranalüüs kinnitas eestikeelse skaala kolmefaktorilist struktuuri sarnaselt esialgsele Rootsi normatiivile: neurootilisus, agressiivsus ja ekstrasvertsus.

Eestikeelse NEO-PI-R neurootilisuse dimensioon korreleerus kõige tugevamalt SSP neurootilisuse faktoriga. NEO-PI-R ekstraversus korreleerus positiivselt SSP alaskaalade seikluste otsimise ja impulsiivsusega ning negatiivselt isoleerumise, psüühilise ärevuse, kehtestavuse puudumise, vastuvõtlikkusega stressile ning umbusklikkusega. NEO-PI-R faktor sotsiaalsus korreleerus positiivselt SSP alaskaala sotsiaalse soovitavusega, negatiivselt agressiivsuse-

ärrituvuse, umbusklikkuse ja kibestumisega. NEO-PI-R faktorid meelekindlus ja avatus olid nõrgemalt seotud SSP-ga. NEO-PI-R meelekindlus korreleerus kõige enam SSP sotsiaalse soovitatavuse ja avatuse faktor seikluste otsimisega. Kokkuvõtvalt, NEO-PI-R ja SSP uurimine kinnitas viimase skaala valiidsust. Erinevused kahe skaala vahel võivad olla tingitud sellest, et SSP on võrreldes NEO-PI-R-ga spetsiifilisema suunaga, püüdes kirjeldada isiksuse patoloogilisi nüansse.

Kokkuvõtvalt näitas SSP eestikeelne versioon kõrget reliaablust ja valiidsust kõrvutatuna Rootsi originaalvalimiga ning ka viiefaktorilise isiksusemudeliga.

Teises uuringus ei olnud olulisi erinevusi patsientide ja kontrollgrupi demograafilistes näitajates. Komorbiidse PH patsientidel oli enam psüühilisi häireid perekonna anamneesis võrrelduna ainult PH patsientidega, samuti oli komorbiidses PH gruppis rohkem suitsetajaid. SSP alaskaalade analüüsis leiti PH patsientide ja tervete kontrollgrupi isiksuse omaduste vahel erinevused. Kõik SSP alaskaalad, välja arvatud isoleerumine ja füüsiline agressioon, olid PH patsientidel oluliselt nihkes võrreldes kontrollgrupiga, sealhulgas seikluste otsimine ja sotsiaalne soovitus madalamad, kõik teised alaskaalad kõrgemad. SSP faktorid neurootilisus ja agressiivsus, kuid mitte ekstraversus, olid oluliselt kõrgemad PH grupis võrreldes kontrollgrupiga.

Komorbiidse paanikahäire ning kontrollgrupi võrdluses leiti oluline erinevus kõigis SSP alaskaalades, välja arvatud seikluste otsimine. Kõik alaskaalad olid komorbiidse PH grupis kõrgemate väärtustega, välja arvatud sotsiaalne soovitus, mis oli madalam. Ainult PH patsiendid demonstreerisid kõrgemat neurootilisust ja agressiivsust, kuid ekstraversus oli sarnane kontrollgrupiga.

Kahe paanikahäirega patsientide grupi võrdluses olid nihked rohkem väljendunud komorbiidse PH grupis, erinevus leiti ärevuse alaskaalade, stressile vastuvõtlikkuse, kibestumise ja ärritavuse skaaladel. Kolmest faktorist erines kahe PH grupi vahel ainult neurootilisus.

Käesoleva paanikahäirega grupis võrreldes remissioonis patsientidega olid oluliselt kõrgemad väärtused ärevuse alaskaaladel ja stressile vastuvõtlikkusel, kuid mitte teistel skaaladel. SSP faktoritest oli ainult neurootilisus oluliselt kõrgem käesoleva PH patsientidel.

Käesoleva PH-ga patsiendid näitasid võrreldes kontrollgrupiga olulist erinevust mitmel SSP alaskaal, välja arvatud impulsiivsus, isoleeritus ja füüsiline agressioon.

Seega võib uuringu tulemusel väita, et paanikahäirega patsiente iseloomustavad väheadaptiivsed isiksusejooned, eriti kõrge neurootilisus ning tugev kalduvus somaatilisele ärevusele.

Kolmandas uuringus leiti kõigi SSP alaskaalade T-skooride langemine 12 nädalase farmakoloogilise ravi järel võrreldes algväärtusega, kuid ükski muutus ei olnud statistiliselt olulisel tasemel. Kõige rohkem vähenesid somaatiline ärevus, vastuvõtlikkus stressile ja psüühiline ärevus.

Leiti mõned nõrgad, statistiliselt mitteolulised seosed SSP alaskaalade ja kliinilise seisundi hinnangu skaalade vahel.

Ükski SSP alaskaalade algväärtusest ei olnud oluliselt seotud ravivastusega. Ainult kõrgem SSP impulsiivsuse tase uuringu alguses oli lähedal statistiliselt olulisele ennustamaks remissiooni mittesaavutamist peale 12 nädalast ravi estsitalopraamiga. Algse impulsiivsuse tase oli statistilisele olulisele lähedaselt seotud ka CGI, HAM-A ja PDSS väiksema langusega raviperioodil.

Järeldused

1. Demonstreerisime, et SSP eestikeelne versioon näitas aktsepteeritavaid psühhomeetrilisi omadusi. SSP mõõdab isiksuse jooni, mis on universaalsed ja haakuvad ühe varasema juhtiva isiksuseteooriaga.
2. SSP suutis lisada asjakohast informatsiooni kliiniliselt olulistele isiksuse dimensioonidele paanikahäirega patsientidel. Me näitasime, et PH patsientide isiksuse struktuur on madala adaptiivsusega. Neid patsiente iseloomustab eelkõige kõrge neurootilisus ja somaatilise ärevuse väljendumine.
3. Enam väljendub isiksusejoonte kõrvalekalle afektiivse komorbiidsusega paanikahäirega patsientidel. Agressiivsus oli seotud agorafobia ja afektiivse komorbiidusega.
4. Kuigi PH patsientide madala adaptiivsusega isiksus näitas kindlat suunda normaliseerumisele 12-nädalase antidepressandi estsitalopraami raviga, ei leidnud me isiksusejoonte olulist seost kliinilise ravitulemusega. Parema farmakoloogilise ravivastusega kaasnes enam väljendunud paranemine isiksuse joontes, mida võib omistada üldiselt paremale enesetunnetusele paranemisel.
5. Võimalikku ennustavat efekti PH ravis antidepressandiga näitas impulsiivsus, mis näib huvipakkuv ning vajab edasist uurimist.

ACKNOWLEDGEMENTS

This work was carried out at the Department of Psychiatry of University of Tartu and at the Psychiatric Clinic of North Estonia Regional Hospital Foundation Psychiatric Clinic.

First, I wanted to express my heartfelt gratitude to my supervisors:

- Eduard Maron MD, PhD inspiring with his excellent scientific advice, support and constructive discussions
- Anu Aluoja, PhD, for her warm guiding support and encouragement. I am grateful for finding always time for consultation during all this long period of work.
- to Innar Tõru MD, PhD for his support

I would like express my gratitude to all the co-authors and colleagues who contributed their time and efforts to this work:

- to Mait Raag MSc for helping me with the statistics and being always helpful.
- to Helina Voogne
- to J. Petter Gustavsson, PhD
- to Jakov Shlik MD, PhD
- to Veiko Vasar MD, PhD
- to study nurses for their correct work Ketlin Veeväli, Birgit Ahas, Jelena Maltseva
- to Mrs Ülle Iher for her time for her help with finishing the manuscript
- to all my colleagues, being supportive and understanding, enable me to organize my everyday work to find a time for my dissertation.
- to the all participants of the studies, being correct and helpful during studies

My gratitude to my family and friends for their support and understanding – especially my husband Vesse and sons for their support

PUBLICATIONS

CURRICULUM VITAE

Full name Ülle Võhma
Date of birth February 23, 1963
Citizenship Estonian
Employment North Estonia Medical Centre Foundation Psychiatric Clinic
Address 52 Paldiski Rd, Tallinn 10614, Estonia
Phone + 372 617 25 00
E-mail ylle.vohma@regionaalhaigla.ee

Education:

1999–2017 University of Tartu, Department of Psychiatry, doctoral student
1996–1999 University of Tartu, Faculty of Medicine, residency in psychiatry
1991–1995 University of Tartu, Faculty of Philosophy, Speciality: Psychology Baccalaureus Scientiarum
1981–1987 University of Tartu, Faculty of Medicine, Department of Paediatrics

Scientific work:

My research interest has been focused on panic disorder and on the association of the disorder with personality traits

Working Experience:

2002– North Estonia Medical Centre Foundation Psychiatric Clinic, Psychiatrist, Senior Doctor, The Head of Department
1999–2002 Tallinn Psychiatric Hospital, Psychiatrist
1988–1996 Tallinn Pelgulinna Hospital Children Outpatient Department, Paediatrician

Publications:

Võhma, Ü.; Raag, M.; Tõru, I.; Aluoja, A.; Maron, E. (2017). Association between personality traits and Escitalopram treatment efficacy in panic disorder. *Nordic Journal of Psychiatry*, 71 (6), 433–440.
Võhma, Ü.; Aluoja, A.; Vasar, V.; Shlik, J.; Maron, E. (2010). Evaluation of personality traits in panic disorder using Swedish universities Scales of Personality. *Journal of Anxiety Disorders*, 24, 141–146.
Aluoja, A.; Voogne, H.; Maron, E.; Gustavsson, J.P.; Võhma, Ü.; Shlik, J. (2009). Personality traits measured by the Swedish universities Scales of Personality: factor structure and position within the five-factor model in an Estonian sample. *Nordic Journal of Psychiatry*, 63, 231–236.

Membership in professional organizations:

Estonian Association of Psychiatrists

ELULOOKIRJELDUS

Nimi: Ülle Võhma
Sünniaeg: 23.02.1963
Kodakondsus: Eesti
Praegune töökoht: SA PER Psühhiaatrikliinik
Aadress: Paldiski maantee 52/6, 10617 Tallinn
Telefon: + 372 617 25 00
E-post: ylle.vohma@regionalhaigla.ee

Haridus:

2009–2017 Tartu Ülikool, arstiteaduskond, psühhiaatrikliinik, doktoriõpe
1996–1999 Tartu Ülikool, psühhiaatria residentuur
1991–1995 Tartu Ülikool filosoofiateaduskond: psühholoogia eriala *baccalaureus scientiarum*
1981–1987 Tartu Riiklik Ülikool, pediaatria eriala

Teadustöö:

uurimistöö on põhiliselt keskendunud paanikahäirele ja selle seostele isiksusejoontega

Töökogemus:

2002– SA Põhja Eesti Regionaalhaigla Psühhiaatrikliinik
psühhiaater, vanemarst, osakonna juhataja
1999–2002 Tallinna Psühhiaatriaigla, psühhiaater
1988–1996 Tallinn Pelgulinna Haigla Kalamaja Lastepolikliinik, pediaater

Publikatsioonid:

Võhma, Ü.; Raag, M.; Tõru, I.; Aluoja, A.; Maron, E. (2017). Paanikahäirega patsientide isikuomaduste seos estsitalopraami ravi efektiivsusega. *Nordic Journal of Psychiatry*. 71 (6): 433–440.
Võhma, Ü.; Aluoja, A.; Vasar, V.; Shlik, J.; Maron, E. (2010). Paanikahäirega patsiendi isikuomaduste hindamine kasutades Rootsi ülikoolide isiksuse-skaalat. *Journal of Anxiety Disorders*, 24, 141–146.
Aluoja, A.; Voogne, H.; Maron, E.; Gustavsson, J.P.; Võhma, Ü.; Shlik, J. (2009). Isiksuse omadused mõõdetuna Rootsi ülikoolide isiksusskaalaga: faktorstruktuur ja positsioon viie-faktorilise isiksuse mudeli suhtes Eesti valimis. *Nordic Journal of Psychiatry*, 63, 231–236.

Kuuluvus erialaühendustesse: Eesti Psühhiaatrite Selts

DISSERTATIONES MEDICINAE UNIVERSITATIS TARTUENSIS

1. **Heidi-Ingrid Maaroo**s. The natural course of gastric ulcer in connection with chronic gastritis and *Helicobacter pylori*. Tartu, 1991.
2. **Mihkel Zilmer**. Na-pump in normal and tumorous brain tissues: Structural, functional and tumorigenesis aspects. Tartu, 1991.
3. **Eero Vasar**. Role of cholecystokinin receptors in the regulation of behaviour and in the action of haloperidol and diazepam. Tartu, 1992.
4. **Tiina Talvik**. Hypoxic-ischaemic brain damage in neonates (clinical, biochemical and brain computed tomographical investigation). Tartu, 1992.
5. **Ants Peetsalu**. Vagotomy in duodenal ulcer disease: A study of gastric acidity, serum pepsinogen I, gastric mucosal histology and *Helicobacter pylori*. Tartu, 1992.
6. **Marika Mikelsaar**. Evaluation of the gastrointestinal microbial ecosystem in health and disease. Tartu, 1992.
7. **Hele Everaus**. Immuno-hormonal interactions in chronic lymphocytic leukaemia and multiple myeloma. Tartu, 1993.
8. **Ruth Mikelsaar**. Etiological factors of diseases in genetically consulted children and newborn screening: dissertation for the commencement of the degree of doctor of medical sciences. Tartu, 1993.
9. **Agu Tamm**. On metabolic action of intestinal microflora: clinical aspects. Tartu, 1993.
10. **Katrin Gross**. Multiple sclerosis in South-Estonia (epidemiological and computed tomographical investigations). Tartu, 1993.
11. **Oivi Uiho**. Childhood coeliac disease in Estonia: occurrence, screening, diagnosis and clinical characterization. Tartu, 1994.
12. **Viiu Tuulik**. The functional disorders of central nervous system of chemistry workers. Tartu, 1994.
13. **Margus Viigimaa**. Primary haemostasis, antiaggregative and anticoagulant treatment of acute myocardial infarction. Tartu, 1994.
14. **Rein Kolk**. Atrial versus ventricular pacing in patients with sick sinus syndrome. Tartu, 1994.
15. **Toomas Podar**. Incidence of childhood onset type 1 diabetes mellitus in Estonia. Tartu, 1994.
16. **Kiira Subi**. The laboratory surveillance of the acute respiratory viral infections in Estonia. Tartu, 1995.
17. **Irja Lutsar**. Infections of the central nervous system in children (epidemiologic, diagnostic and therapeutic aspects, long term outcome). Tartu, 1995.
18. **Aavo Lang**. The role of dopamine, 5-hydroxytryptamine, sigma and NMDA receptors in the action of antipsychotic drugs. Tartu, 1995.
19. **Andrus Arak**. Factors influencing the survival of patients after radical surgery for gastric cancer. Tartu, 1996.

20. **Tõnis Karki.** Quantitative composition of the human lactoflora and method for its examination. Tartu, 1996.
21. **Reet Mändar.** Vaginal microflora during pregnancy and its transmission to newborn. Tartu, 1996.
22. **Triin Remmel.** Primary biliary cirrhosis in Estonia: epidemiology, clinical characterization and prognostication of the course of the disease. Tartu, 1996.
23. **Toomas Kivastik.** Mechanisms of drug addiction: focus on positive reinforcing properties of morphine. Tartu, 1996.
24. **Paavo Pokk.** Stress due to sleep deprivation: focus on GABA_A receptor-chloride ionophore complex. Tartu, 1996.
25. **Kristina Allikmets.** Renin system activity in essential hypertension. Associations with atherothrombogenic cardiovascular risk factors and with the efficacy of calcium antagonist treatment. Tartu, 1996.
26. **Triin Parik.** Oxidative stress in essential hypertension: Associations with metabolic disturbances and the effects of calcium antagonist treatment. Tartu, 1996.
27. **Svetlana Päi.** Factors promoting heterogeneity of the course of rheumatoid arthritis. Tartu, 1997.
28. **Maarike Sallo.** Studies on habitual physical activity and aerobic fitness in 4 to 10 years old children. Tartu, 1997.
29. **Paul Naaber.** *Clostridium difficile* infection and intestinal microbial ecology. Tartu, 1997.
30. **Rein Pähkla.** Studies in pinoline pharmacology. Tartu, 1997.
31. **Andrus Juhan Voitk.** Outpatient laparoscopic cholecystectomy. Tartu, 1997.
32. **Joel Starkopf.** Oxidative stress and ischaemia-reperfusion of the heart. Tartu, 1997.
33. **Janika Kõrv.** Incidence, case-fatality and outcome of stroke. Tartu, 1998.
34. **Ülla Linnamägi.** Changes in local cerebral blood flow and lipid peroxidation following lead exposure in experiment. Tartu, 1998.
35. **Ave Minajeva.** Sarcoplasmic reticulum function: comparison of atrial and ventricular myocardium. Tartu, 1998.
36. **Oleg Milenin.** Reconstruction of cervical part of esophagus by revascularised ileal autografts in dogs. A new complex multistage method. Tartu, 1998.
37. **Sergei Pakriev.** Prevalence of depression, harmful use of alcohol and alcohol dependence among rural population in Udmurtia. Tartu, 1998.
38. **Allen Kaasik.** Thyroid hormone control over β -adrenergic signalling system in rat atria. Tartu, 1998.
39. **Vallo Matto.** Pharmacological studies on anxiogenic and antiaggressive properties of antidepressants. Tartu, 1998.
40. **Maire Vasar.** Allergic diseases and bronchial hyperreactivity in Estonian children in relation to environmental influences. Tartu, 1998.
41. **Kaja Julge.** Humoral immune responses to allergens in early childhood. Tartu, 1998.

42. **Heli Grünberg.** The cardiovascular risk of Estonian schoolchildren. A cross-sectional study of 9-, 12- and 15-year-old children. Tartu, 1998.
43. **Epp Sepp.** Formation of intestinal microbial ecosystem in children. Tartu, 1998.
44. **Mai Ots.** Characteristics of the progression of human and experimental glomerulopathies. Tartu, 1998.
45. **Tiina Ristimäe.** Heart rate variability in patients with coronary artery disease. Tartu, 1998.
46. **Leho Kõiv.** Reaction of the sympatho-adrenal and hypothalamo-pituitary-adrenocortical system in the acute stage of head injury. Tartu, 1998.
47. **Bela Adojaan.** Immune and genetic factors of childhood onset IDDM in Estonia. An epidemiological study. Tartu, 1999.
48. **Jakov Shlik.** Psychophysiological effects of cholecystokinin in humans. Tartu, 1999.
49. **Kai Kisand.** Autoantibodies against dehydrogenases of α -ketoacids. Tartu, 1999.
50. **Toomas Marandi.** Drug treatment of depression in Estonia. Tartu, 1999.
51. **Ants Kask.** Behavioural studies on neuropeptide Y. Tartu, 1999.
52. **Ello-Rahel Karelson.** Modulation of adenylate cyclase activity in the rat hippocampus by neuropeptide galanin and its chimeric analogs. Tartu, 1999.
53. **Tanel Laisaar.** Treatment of pleural empyema — special reference to intrapleural therapy with streptokinase and surgical treatment modalities. Tartu, 1999.
54. **Eve Pihl.** Cardiovascular risk factors in middle-aged former athletes. Tartu, 1999.
55. **Katrin Õunap.** Phenylketonuria in Estonia: incidence, newborn screening, diagnosis, clinical characterization and genotype/phenotype correlation. Tartu, 1999.
56. **Siiri Kõljalg.** *Acinetobacter* – an important nosocomial pathogen. Tartu, 1999.
57. **Helle Karro.** Reproductive health and pregnancy outcome in Estonia: association with different factors. Tartu, 1999.
58. **Heili Varendi.** Behavioral effects observed in human newborns during exposure to naturally occurring odors. Tartu, 1999.
59. **Anneli Beilmann.** Epidemiology of epilepsy in children and adolescents in Estonia. Prevalence, incidence, and clinical characteristics. Tartu, 1999.
60. **Vallo Volke.** Pharmacological and biochemical studies on nitric oxide in the regulation of behaviour. Tartu, 1999.
61. **Pilvi Ilves.** Hypoxic-ischaemic encephalopathy in asphyxiated term infants. A prospective clinical, biochemical, ultrasonographical study. Tartu, 1999.
62. **Anti Kalda.** Oxygen-glucose deprivation-induced neuronal death and its pharmacological prevention in cerebellar granule cells. Tartu, 1999.
63. **Eve-Irene Lepist.** Oral peptide prodrugs – studies on stability and absorption. Tartu, 2000.

64. **Jana Kivastik.** Lung function in Estonian schoolchildren: relationship with anthropometric indices and respiratory symptoms, reference values for dynamic spirometry. Tartu, 2000.
65. **Karin Kull.** Inflammatory bowel disease: an immunogenetic study. Tartu, 2000.
66. **Kaire Innos.** Epidemiological resources in Estonia: data sources, their quality and feasibility of cohort studies. Tartu, 2000.
67. **Tamara Vorobjova.** Immune response to *Helicobacter pylori* and its association with dynamics of chronic gastritis and epithelial cell turnover in antrum and corpus. Tartu, 2001.
68. **Ruth Kalda.** Structure and outcome of family practice quality in the changing health care system of Estonia. Tartu, 2001.
69. **Annika Krüüner.** *Mycobacterium tuberculosis* – spread and drug resistance in Estonia. Tartu, 2001.
70. **Marlit Veldi.** Obstructive Sleep Apnoea: Computerized Endopharyngeal Myotonometry of the Soft Palate and Lingual Musculature. Tartu, 2001.
71. **Anneli Uusküla.** Epidemiology of sexually transmitted diseases in Estonia in 1990–2000. Tartu, 2001.
72. **Ade Kallas.** Characterization of antibodies to coagulation factor VIII. Tartu, 2002.
73. **Heidi Annuk.** Selection of medicinal plants and intestinal lactobacilli as antimicrobial components for functional foods. Tartu, 2002.
74. **Aet Lukmann.** Early rehabilitation of patients with ischaemic heart disease after surgical revascularization of the myocardium: assessment of health-related quality of life, cardiopulmonary reserve and oxidative stress. A clinical study. Tartu, 2002.
75. **Maigi Eisen.** Pathogenesis of Contact Dermatitis: participation of Oxidative Stress. A clinical – biochemical study. Tartu, 2002.
76. **Piret Hussar.** Histology of the post-traumatic bone repair in rats. Elaboration and use of a new standardized experimental model – bicortical perforation of tibia compared to internal fracture and resection osteotomy. Tartu, 2002.
77. **Tõnu Rätsep.** Aneurysmal subarachnoid haemorrhage: Noninvasive monitoring of cerebral haemodynamics. Tartu, 2002.
78. **Marju Herodes.** Quality of life of people with epilepsy in Estonia. Tartu, 2003.
79. **Katre Maasalu.** Changes in bone quality due to age and genetic disorders and their clinical expressions in Estonia. Tartu, 2003.
80. **Toomas Sillakivi.** Perforated peptic ulcer in Estonia: epidemiology, risk factors and relations with *Helicobacter pylori*. Tartu, 2003.
81. **Leena Puksa.** Late responses in motor nerve conduction studies. F and A waves in normal subjects and patients with neuropathies. Tartu, 2003.
82. **Krista Lõivukene.** *Helicobacter pylori* in gastric microbial ecology and its antimicrobial susceptibility pattern. Tartu, 2003.

83. **Helgi Kolk.** Dyspepsia and *Helicobacter pylori* infection: the diagnostic value of symptoms, treatment and follow-up of patients referred for upper gastrointestinal endoscopy by family physicians. Tartu, 2003.
84. **Helena Soomer.** Validation of identification and age estimation methods in forensic odontology. Tartu, 2003.
85. **Kersti Oselin.** Studies on the human MDR1, MRP1, and MRP2 ABC transporters: functional relevance of the genetic polymorphisms in the *MDR1* and *MRP1* gene. Tartu, 2003.
86. **Jaan Soplemann.** Peptic ulcer haemorrhage in Estonia: epidemiology, prognostic factors, treatment and outcome. Tartu, 2003.
87. **Margot Peetsalu.** Long-term follow-up after vagotomy in duodenal ulcer disease: recurrent ulcer, changes in the function, morphology and *Helicobacter pylori* colonisation of the gastric mucosa. Tartu, 2003.
88. **Kersti Klaamas.** Humoral immune response to *Helicobacter pylori* a study of host-dependent and microbial factors. Tartu, 2003.
89. **Pille Taba.** Epidemiology of Parkinson's disease in Tartu, Estonia. Prevalence, incidence, clinical characteristics, and pharmacoepidemiology. Tartu, 2003.
90. **Alar Veraksitš.** Characterization of behavioural and biochemical phenotype of cholecystokinin-2 receptor deficient mice: changes in the function of the dopamine and endopioidergic system. Tartu, 2003.
91. **Ingrid Kalev.** CC-chemokine receptor 5 (CCR5) gene polymorphism in Estonians and in patients with Type I and Type II diabetes mellitus. Tartu, 2003.
92. **Lumme Kadaja.** Molecular approach to the regulation of mitochondrial function in oxidative muscle cells. Tartu, 2003.
93. **Aive Liigant.** Epidemiology of primary central nervous system tumours in Estonia from 1986 to 1996. Clinical characteristics, incidence, survival and prognostic factors. Tartu, 2004.
94. **Andres, Kulla.** Molecular characteristics of mesenchymal stroma in human astrocytic gliomas. Tartu, 2004.
95. **Mari Järvelaid.** Health damaging risk behaviours in adolescence. Tartu, 2004.
96. **Ülle Pechter.** Progression prevention strategies in chronic renal failure and hypertension. An experimental and clinical study. Tartu, 2004.
97. **Gunnar Tasa.** Polymorphic glutathione S-transferases – biology and role in modifying genetic susceptibility to senile cataract and primary open angle glaucoma. Tartu, 2004.
98. **Tuuli Käämbre.** Intracellular energetic unit: structural and functional aspects. Tartu, 2004.
99. **Vitali Vassiljev.** Influence of nitric oxide synthase inhibitors on the effects of ethanol after acute and chronic ethanol administration and withdrawal. Tartu, 2004.

100. **Aune Rehema.** Assessment of nonhaem ferrous iron and glutathione redox ratio as markers of pathogeneticity of oxidative stress in different clinical groups. Tartu, 2004.
101. **Evelin Seppet.** Interaction of mitochondria and ATPases in oxidative muscle cells in normal and pathological conditions. Tartu, 2004.
102. **Eduard Maron.** Serotonin function in panic disorder: from clinical experiments to brain imaging and genetics. Tartu, 2004.
103. **Marje Oona.** *Helicobacter pylori* infection in children: epidemiological and therapeutic aspects. Tartu, 2004.
104. **Kersti Kokk.** Regulation of active and passive molecular transport in the testis. Tartu, 2005.
105. **Vladimir Järv.** Cross-sectional imaging for pretreatment evaluation and follow-up of pelvic malignant tumours. Tartu, 2005.
106. **Andre Õun.** Epidemiology of adult epilepsy in Tartu, Estonia. Incidence, prevalence and medical treatment. Tartu, 2005.
107. **Piibe Muda.** Homocysteine and hypertension: associations between homocysteine and essential hypertension in treated and untreated hypertensive patients with and without coronary artery disease. Tartu, 2005.
108. **Küllü Kingo.** The interleukin-10 family cytokines gene polymorphisms in plaque psoriasis. Tartu, 2005.
109. **Mati Merila.** Anatomy and clinical relevance of the glenohumeral joint capsule and ligaments. Tartu, 2005.
110. **Epp Songisepp.** Evaluation of technological and functional properties of the new probiotic *Lactobacillus fermentum* ME-3. Tartu, 2005.
111. **Tiia Ainla.** Acute myocardial infarction in Estonia: clinical characteristics, management and outcome. Tartu, 2005.
112. **Andres Sell.** Determining the minimum local anaesthetic requirements for hip replacement surgery under spinal anaesthesia – a study employing a spinal catheter. Tartu, 2005.
113. **Tiia Tamme.** Epidemiology of odontogenic tumours in Estonia. Pathogenesis and clinical behaviour of ameloblastoma. Tartu, 2005.
114. **Triine Annus.** Allergy in Estonian schoolchildren: time trends and characteristics. Tartu, 2005.
115. **Tiia Voor.** Microorganisms in infancy and development of allergy: comparison of Estonian and Swedish children. Tartu, 2005.
116. **Priit Kasenõmm.** Indicators for tonsillectomy in adults with recurrent tonsillitis – clinical, microbiological and pathomorphological investigations. Tartu, 2005.
117. **Eva Zusinaite.** Hepatitis C virus: genotype identification and interactions between viral proteases. Tartu, 2005.
118. **Piret Kõll.** Oral lactoflora in chronic periodontitis and periodontal health. Tartu, 2006.
119. **Tiina Stelmach.** Epidemiology of cerebral palsy and unfavourable neurodevelopmental outcome in child population of Tartu city and county, Estonia Prevalence, clinical features and risk factors. Tartu, 2006.

120. **Katrin Pudersell.** Tropane alkaloid production and riboflavine excretion in the field and tissue cultures of henbane (*Hyoscyamus niger* L.). Tartu, 2006.
121. **Küllil Jaako.** Studies on the role of neurogenesis in brain plasticity. Tartu, 2006.
122. **Aare Märtsen.** Lower limb lengthening: experimental studies of bone regeneration and long-term clinical results. Tartu, 2006.
123. **Heli Tähepõld.** Patient consultation in family medicine. Tartu, 2006.
124. **Stanislav Liskmann.** Peri-implant disease: pathogenesis, diagnosis and treatment in view of both inflammation and oxidative stress profiling. Tartu, 2006.
125. **Ruth Rudissaar.** Neuropharmacology of atypical antipsychotics and an animal model of psychosis. Tartu, 2006.
126. **Helena Andreson.** Diversity of *Helicobacter pylori* genotypes in Estonian patients with chronic inflammatory gastric diseases. Tartu, 2006.
127. **Katrin Pruus.** Mechanism of action of antidepressants: aspects of serotonergic system and its interaction with glutamate. Tartu, 2006.
128. **Priit Põder.** Clinical and experimental investigation: relationship of ischaemia/reperfusion injury with oxidative stress in abdominal aortic aneurysm repair and in extracranial brain artery endarterectomy and possibilities of protection against ischaemia using a glutathione analogue in a rat model of global brain ischaemia. Tartu, 2006.
129. **Marika Tammaru.** Patient-reported outcome measurement in rheumatoid arthritis. Tartu, 2006.
130. **Tiia Reimand.** Down syndrome in Estonia. Tartu, 2006.
131. **Diva Eensoo.** Risk-taking in traffic and Markers of Risk-Taking Behaviour in Schoolchildren and Car Drivers. Tartu, 2007.
132. **Riina Vibo.** The third stroke registry in Tartu, Estonia from 2001 to 2003: incidence, case-fatality, risk factors and long-term outcome. Tartu, 2007.
133. **Chris Pruunsild.** Juvenile idiopathic arthritis in children in Estonia. Tartu, 2007.
134. **Eve Õiglane-Šlik.** Angelman and Prader-Willi syndromes in Estonia. Tartu, 2007.
135. **Kadri Haller.** Antibodies to follicle stimulating hormone. Significance in female infertility. Tartu, 2007.
136. **Pille Ööpik.** Management of depression in family medicine. Tartu, 2007.
137. **Jaak Kals.** Endothelial function and arterial stiffness in patients with atherosclerosis and in healthy subjects. Tartu, 2007.
138. **Priit Kampus.** Impact of inflammation, oxidative stress and age on arterial stiffness and carotid artery intima-media thickness. Tartu, 2007.
139. **Margus Punab.** Male fertility and its risk factors in Estonia. Tartu, 2007.
140. **Alar Toom.** Heterotopic ossification after total hip arthroplasty: clinical and pathogenetic investigation. Tartu, 2007.

141. **Lea Pehme.** Epidemiology of tuberculosis in Estonia 1991–2003 with special regard to extrapulmonary tuberculosis and delay in diagnosis of pulmonary tuberculosis. Tartu, 2007.
142. **Juri Karjagin.** The pharmacokinetics of metronidazole and meropenem in septic shock. Tartu, 2007.
143. **Inga Talvik.** Inflicted traumatic brain injury shaken baby syndrome in Estonia – epidemiology and outcome. Tartu, 2007.
144. **Tarvo Rajasalu.** Autoimmune diabetes: an immunological study of type 1 diabetes in humans and in a model of experimental diabetes (in RIP-B7.1 mice). Tartu, 2007.
145. **Inga Karu.** Ischaemia-reperfusion injury of the heart during coronary surgery: a clinical study investigating the effect of hyperoxia. Tartu, 2007.
146. **Peeter Padrik.** Renal cell carcinoma: Changes in natural history and treatment of metastatic disease. Tartu, 2007.
147. **Neve Vendt.** Iron deficiency and iron deficiency anaemia in infants aged 9 to 12 months in Estonia. Tartu, 2008.
148. **Lenne-Triin Heidmets.** The effects of neurotoxins on brain plasticity: focus on neural Cell Adhesion Molecule. Tartu, 2008.
149. **Paul Korrovits.** Asymptomatic inflammatory prostatitis: prevalence, etiological factors, diagnostic tools. Tartu, 2008.
150. **Annika Reintam.** Gastrointestinal failure in intensive care patients. Tartu, 2008.
151. **Kristiina Roots.** Cationic regulation of Na-pump in the normal, Alzheimer's and CCK₂ receptor-deficient brain. Tartu, 2008.
152. **Helen Puusepp.** The genetic causes of mental retardation in Estonia: fragile X syndrome and creatine transporter defect. Tartu, 2009.
153. **Kristiina Rull.** Human chorionic gonadotropin beta genes and recurrent miscarriage: expression and variation study. Tartu, 2009.
154. **Margus Eimre.** Organization of energy transfer and feedback regulation in oxidative muscle cells. Tartu, 2009.
155. **Maire Link.** Transcription factors FoxP3 and AIRE: autoantibody associations. Tartu, 2009.
156. **Kai Haldre.** Sexual health and behaviour of young women in Estonia. Tartu, 2009.
157. **Kaur Liivak.** Classical form of congenital adrenal hyperplasia due to 21-hydroxylase deficiency in Estonia: incidence, genotype and phenotype with special attention to short-term growth and 24-hour blood pressure. Tartu, 2009.
158. **Kersti Ehrlich.** Antioxidative glutathione analogues (UPF peptides) – molecular design, structure-activity relationships and testing the protective properties. Tartu, 2009.
159. **Anneli Rätsep.** Type 2 diabetes care in family medicine. Tartu, 2009.
160. **Silver Türk.** Etiopathogenetic aspects of chronic prostatitis: role of mycoplasmas, coryneform bacteria and oxidative stress. Tartu, 2009.

161. **Kaire Heilman**. Risk markers for cardiovascular disease and low bone mineral density in children with type 1 diabetes. Tartu, 2009.
162. **Kristi Rüütel**. HIV-epidemic in Estonia: injecting drug use and quality of life of people living with HIV. Tartu, 2009.
163. **Triin Eller**. Immune markers in major depression and in antidepressive treatment. Tartu, 2009.
164. **Siim Suutre**. The role of TGF- β isoforms and osteoprogenitor cells in the pathogenesis of heterotopic ossification. An experimental and clinical study of hip arthroplasty. Tartu, 2010.
165. **Kai Kliiman**. Highly drug-resistant tuberculosis in Estonia: Risk factors and predictors of poor treatment outcome. Tartu, 2010.
166. **Inga Villa**. Cardiovascular health-related nutrition, physical activity and fitness in Estonia. Tartu, 2010.
167. **Tõnis Org**. Molecular function of the first PHD finger domain of Auto-immune Regulator protein. Tartu, 2010.
168. **Tuuli Metsvaht**. Optimal antibacterial therapy of neonates at risk of early onset sepsis. Tartu, 2010.
169. **Jaanus Kahu**. Kidney transplantation: Studies on donor risk factors and mycophenolate mofetil. Tartu, 2010.
170. **Koit Reimand**. Autoimmunity in reproductive failure: A study on associated autoantibodies and autoantigens. Tartu, 2010.
171. **Mart Kull**. Impact of vitamin D and hypolactasia on bone mineral density: a population based study in Estonia. Tartu, 2010.
172. **Rael Laugesaar**. Stroke in children – epidemiology and risk factors. Tartu, 2010.
173. **Mark Braschinsky**. Epidemiology and quality of life issues of hereditary spastic paraplegia in Estonia and implementation of genetic analysis in everyday neurologic practice. Tartu, 2010.
174. **Kadri Suija**. Major depression in family medicine: associated factors, recurrence and possible intervention. Tartu, 2010.
175. **Jarno Habicht**. Health care utilisation in Estonia: socioeconomic determinants and financial burden of out-of-pocket payments. Tartu, 2010.
176. **Kristi Abram**. The prevalence and risk factors of rosacea. Subjective disease perception of rosacea patients. Tartu, 2010.
177. **Malle Kuum**. Mitochondrial and endoplasmic reticulum cation fluxes: Novel roles in cellular physiology. Tartu, 2010.
178. **Rita Teek**. The genetic causes of early onset hearing loss in Estonian children. Tartu, 2010.
179. **Daisy Volmer**. The development of community pharmacy services in Estonia – public and professional perceptions 1993–2006. Tartu, 2010.
180. **Jelena Lissitsina**. Cytogenetic causes in male infertility. Tartu, 2011.
181. **Delia Lepik**. Comparison of gunshot injuries caused from Tokarev, Makarov and Glock 19 pistols at different firing distances. Tartu, 2011.
182. **Ene-Renate Pähkla**. Factors related to the efficiency of treatment of advanced periodontitis. Tartu, 2011.

183. **Maarja Krass.** L-Arginine pathways and antidepressant action. Tartu, 2011.
184. **Taavi Lai.** Population health measures to support evidence-based health policy in Estonia. Tartu, 2011.
185. **Tiit Salum.** Similarity and difference of temperature-dependence of the brain sodium pump in normal, different neuropathological, and aberrant conditions and its possible reasons. Tartu, 2011.
186. **Tõnu Vooder.** Molecular differences and similarities between histological subtypes of non-small cell lung cancer. Tartu, 2011.
187. **Jelena Štšepetova.** The characterisation of intestinal lactic acid bacteria using bacteriological, biochemical and molecular approaches. Tartu, 2011.
188. **Radko Avi.** Natural polymorphisms and transmitted drug resistance in Estonian HIV-1 CRF06_cpx and its recombinant viruses. Tartu, 2011, 116 p.
189. **Edward Laane.** Multiparameter flow cytometry in haematological malignancies. Tartu, 2011, 152 p.
190. **Triin Jagomägi.** A study of the genetic etiology of nonsyndromic cleft lip and palate. Tartu, 2011, 158 p.
191. **Ivo Laidmäe.** Fibrin glue of fish (*Salmo salar*) origin: immunological study and development of new pharmaceutical preparation. Tartu, 2012, 150 p.
192. **Ülle Parm.** Early mucosal colonisation and its role in prediction of invasive infection in neonates at risk of early onset sepsis. Tartu, 2012, 168 p.
193. **Kaupo Teesalu.** Autoantibodies against desmin and transglutaminase 2 in celiac disease: diagnostic and functional significance. Tartu, 2012, 142 p.
194. **Maksim Zagura.** Biochemical, functional and structural profiling of arterial damage in atherosclerosis. Tartu, 2012, 162 p.
195. **Vivian Kont.** Autoimmune regulator: characterization of thymic gene regulation and promoter methylation. Tartu, 2012, 134 p.
196. **Pirje Hütt.** Functional properties, persistence, safety and efficacy of potential probiotic lactobacilli. Tartu, 2012, 246 p.
197. **Innar Tõru.** Serotonergic modulation of CCK-4- induced panic. Tartu, 2012, 132 p.
198. **Sigrid Vorobjov.** Drug use, related risk behaviour and harm reduction interventions utilization among injecting drug users in Estonia: implications for drug policy. Tartu, 2012, 120 p.
199. **Martin Serg.** Therapeutic aspects of central haemodynamics, arterial stiffness and oxidative stress in hypertension. Tartu, 2012, 156 p.
200. **Jaanika Kumm.** Molecular markers of articular tissues in early knee osteoarthritis: a population-based longitudinal study in middle-aged subjects. Tartu, 2012, 159 p.
201. **Kertu Rünkorg.** Functional changes of dopamine, endopioid and endocannabinoid systems in CCK2 receptor deficient mice. Tartu, 2012, 125 p.
202. **Mai Blöndal.** Changes in the baseline characteristics, management and outcomes of acute myocardial infarction in Estonia. Tartu, 2012, 127 p.

203. **Jana Lass.** Epidemiological and clinical aspects of medicines use in children in Estonia. Tartu, 2012, 170 p.
204. **Kai Truusalu.** Probiotic lactobacilli in experimental persistent *Salmonella* infection. Tartu, 2013, 139 p.
205. **Oksana Jagur.** Temporomandibular joint diagnostic imaging in relation to pain and bone characteristics. Long-term results of arthroscopic treatment. Tartu, 2013, 126 p.
206. **Katrin Sikk.** Manganese-ephedrone intoxication – pathogenesis of neurological damage and clinical symptomatology. Tartu, 2013, 125 p.
207. **Kai Blöndal.** Tuberculosis in Estonia with special emphasis on drug-resistant tuberculosis: Notification rate, disease recurrence and mortality. Tartu, 2013, 151 p.
208. **Marju Puurand.** Oxidative phosphorylation in different diseases of gastric mucosa. Tartu, 2013, 123 p.
209. **Aili Tagoma.** Immune activation in female infertility: Significance of autoantibodies and inflammatory mediators. Tartu, 2013, 135 p.
210. **Liis Sabre.** Epidemiology of traumatic spinal cord injury in Estonia. Brain activation in the acute phase of traumatic spinal cord injury. Tartu, 2013, 135 p.
211. **Merit Lamp.** Genetic susceptibility factors in endometriosis. Tartu, 2013, 125 p.
212. **Erik Salum.** Beneficial effects of vitamin D and angiotensin II receptor blocker on arterial damage. Tartu, 2013, 167 p.
213. **Maire Karelson.** Viteligo: clinical aspects, quality of life and the role of melanocortin system in pathogenesis. Tartu, 2013, 153 p.
214. **Kuldar Kaljurand.** Prevalence of exfoliation syndrome in Estonia and its clinical significance. Tartu, 2013, 113 p.
215. **Raido Paasma.** Clinical study of methanol poisoning: handling large outbreaks, treatment with antidotes, and long-term outcomes. Tartu, 2013, 96 p.
216. **Anne Kleinberg.** Major depression in Estonia: prevalence, associated factors, and use of health services. Tartu, 2013, 129 p.
217. **Triin Eglit.** Obesity, impaired glucose regulation, metabolic syndrome and their associations with high-molecular-weight adiponectin levels. Tartu, 2014, 115 p.
218. **Kristo Ausmees.** Reproductive function in middle-aged males: Associations with prostate, lifestyle and couple infertility status. Tartu, 2014, 125 p.
219. **Kristi Huik.** The influence of host genetic factors on the susceptibility to HIV and HCV infections among intravenous drug users. Tartu, 2014, 144 p.
220. **Liina Tserel.** Epigenetic profiles of monocytes, monocyte-derived macrophages and dendritic cells. Tartu, 2014, 143 p.
221. **Irina Kerna.** The contribution of *ADAM12* and *CILP* genes to the development of knee osteoarthritis. Tartu, 2014, 152 p.

222. **Ingrid Liiv.** Autoimmune regulator protein interaction with DNA-dependent protein kinase and its role in apoptosis. Tartu, 2014, 143 p.
223. **Liivi Maddison.** Tissue perfusion and metabolism during intra-abdominal hypertension. Tartu, 2014, 103 p.
224. **Krista Ress.** Childhood coeliac disease in Estonia, prevalence in atopic dermatitis and immunological characterisation of coexistence. Tartu, 2014, 124 p.
225. **Kai Muru.** Prenatal screening strategies, long-term outcome of children with marked changes in maternal screening tests and the most common syndromic heart anomalies in Estonia. Tartu, 2014, 189 p.
226. **Kaja Rahu.** Morbidity and mortality among Baltic Chernobyl cleanup workers: a register-based cohort study. Tartu, 2014, 155 p.
227. **Klari Noormets.** The development of diabetes mellitus, fertility and energy metabolism disturbances in a Wfs1-deficient mouse model of Wolfram syndrome. Tartu, 2014, 132 p.
228. **Liis Toome.** Very low gestational age infants in Estonia. Tartu, 2014, 183 p.
229. **Ceith Nikkolo.** Impact of different mesh parameters on chronic pain and foreign body feeling after open inguinal hernia repair. Tartu, 2014, 132 p.
230. **Vadim Brjalin.** Chronic hepatitis C: predictors of treatment response in Estonian patients. Tartu, 2014, 122 p.
231. **Vahur Metsna.** Anterior knee pain in patients following total knee arthroplasty: the prevalence, correlation with patellar cartilage impairment and aspects of patellofemoral congruence. Tartu, 2014, 130 p.
232. **Marju Kase.** Glioblastoma multiforme: possibilities to improve treatment efficacy. Tartu, 2015, 137 p.
233. **Riina Runnel.** Oral health among elementary school children and the effects of polyol candies on the prevention of dental caries. Tartu, 2015, 112 p.
234. **Made Laanpere.** Factors influencing women's sexual health and reproductive choices in Estonia. Tartu, 2015, 176 p.
235. **Andres Lust.** Water mediated solid state transformations of a polymorphic drug – effect on pharmaceutical product performance. Tartu, 2015, 134 p.
236. **Anna Klugman.** Functionality related characterization of pretreated wood lignin, cellulose and polyvinylpyrrolidone for pharmaceutical applications. Tartu, 2015, 156 p.
237. **Triin Laisk-Podar.** Genetic variation as a modulator of susceptibility to female infertility and a source for potential biomarkers. Tartu, 2015, 155 p.
238. **Mailis Tõnisson.** Clinical picture and biochemical changes in blood in children with acute alcohol intoxication. Tartu, 2015, 100 p.
239. **Kadri Tamme.** High volume haemodiafiltration in treatment of severe sepsis – impact on pharmacokinetics of antibiotics and inflammatory response. Tartu, 2015, 133 p.

240. **Kai Part.** Sexual health of young people in Estonia in a social context: the role of school-based sexuality education and youth-friendly counseling services. Tartu, 2015, 203 p.
241. **Urve Paaver.** New perspectives for the amorphization and physical stabilization of poorly water-soluble drugs and understanding their dissolution behavior. Tartu, 2015, 139 p.
242. **Aleksandr Peet.** Intrauterine and postnatal growth in children with HLA-conferred susceptibility to type 1 diabetes. Tartu. 2015, 146 p.
243. **Piret Mitt.** Healthcare-associated infections in Estonia – epidemiology and surveillance of bloodstream and surgical site infections. Tartu, 2015, 145 p.
244. **Merli Saare.** Molecular Profiling of Endometriotic Lesions and Endometriosis of Endometriosis Patients. Tartu, 2016, 129 p.
245. **Kaja-Triin Laisaar.** People living with HIV in Estonia: Engagement in medical care and methods of increasing adherence to antiretroviral therapy and safe sexual behavior. Tartu, 2016, 132 p.
246. **Eero Merilind.** Primary health care performance: impact of payment and practice-based characteristics. Tartu, 2016, 120 p.
247. **Jaanika Kärner.** Cytokine-specific autoantibodies in AIRE deficiency. Tartu, 2016, 182 p.
248. **Kaido Paapstel.** Metabolomic profile of arterial stiffness and early biomarkers of renal damage in atherosclerosis. Tartu, 2016, 173 p.
249. **Liidia Kiisk.** Long-term nutritional study: anthropometrical and clinico-laboratory assessments in renal replacement therapy patients after intensive nutritional counselling. Tartu, 2016, 207 p.
250. **Georgi Nellis.** The use of excipients in medicines administered to neonates in Europe. Tartu, 2017, 159 p.
251. **Aleksei Rakitin.** Metabolic effects of acute and chronic treatment with valproic acid in people with epilepsy. Tartu, 2017, 125 p.
252. **Eveli Kallas.** The influence of immunological markers to susceptibility to HIV, HBV, and HCV infections among persons who inject drugs. Tartu, 2017, 138 p.
253. **Tiina Freimann.** Musculoskeletal pain among nurses: prevalence, risk factors, and intervention. Tartu, 2017, 125 p.
254. **Evelyn Aaviksoo.** Sickness absence in Estonia: determinants and influence of the sick-pay cut reform. Tartu, 2017, 121 p.
255. **Kalev Nõupuu.** Autosomal-recessive Stargardt disease: phenotypic heterogeneity and genotype-phenotype associations. Tartu, 2017, 131 p.
256. **Ho Duy Binh.** Osteogenesis imperfecta in Vietnam. Tartu, 2017, 125 p.
257. **Uku Haljasorg.** Transcriptional mechanisms in thymic central tolerance. Tartu, 2017, 147 p.
258. **Živile Riispere.** IgA Nephropathy study according to the Oxford Classification: IgA Nephropathy clinical-morphological correlations, disease progression and the effect of renoprotective therapy. Tartu, 2017, 129 p.

259. **Hiie Soeorg**. Coagulase-negative staphylococci in gut of preterm neonates and in breast milk of their mothers. Tartu, 2017, 216 p.
260. **Anne-Mari Anton Willmore**. Silver nanoparticles for cancer research. Tartu, 2017, 132 p.
261. **Ott Laius**. Utilization of osteoporosis medicines, medication adherence and the trend in osteoporosis related hip fractures in Estonia. Tartu, 2017, 134 p.
262. **Alar Aab**. Insights into molecular mechanisms of asthma and atopic dermatitis. Tartu, 2017, 164 p.
263. **Sander Pajusalu**. Genome-wide diagnostics of Mendelian disorders: from chromosomal microarrays to next-generation sequencing. Tartu, 2017, 146 p.
264. **Mikk Jürisson**. Health and economic impact of hip fracture in Estonia. Tartu, 2017, 164 p.
265. **Kaspar Tootsi**. Cardiovascular and metabolomic profiling of osteoarthritis. Tartu, 2017, 150 p.
266. **Mario Saare**. The influence of AIRE on gene expression – studies of transcriptional regulatory mechanisms in cell culture systems. Tartu, 2017, 172 p.
267. **Piia Jõgi**. Epidemiological and clinical characteristics of pertussis in Estonia. Tartu, 2018, 168 p.
268. **Elle Põldoja**. Structure and blood supply of the superior part of the shoulder joint capsule. Tartu, 2018, 116 p.
269. **Minh Son Nguyen**. Oral health status and prevalence of temporomandibular disorders in 65–74-year-olds in Vietnam. Tartu, 2018, 182 p.
270. **Kristian Semjonov**. Development of pharmaceutical quench-cooled molten and melt-electrospun solid dispersions for poorly water-soluble indomethacin. Tartu, 2018, 125 p.
271. **Janne Tiigimäe-Saar**. Botulinum neurotoxin type A treatment for sialorrhea in central nervous system diseases. Tartu, 2018, 109 p.
272. **Veiko Vengerfeldt**. Apical periodontitis: prevalence and etiopathogenetic aspects. Tartu, 2018, 150 p.
273. **Rudolf Bichele**. TNF superfamily and AIRE at the crossroads of thymic differentiation and host protection against *Candida albicans* infection. Tartu, 2018, 153 p.
274. **Olga Tšuiiko**. Unravelling Chromosomal Instability in Mammalian Pre-implantation Embryos Using Single-Cell Genomics. Tartu, 2018, 169 p.
275. **Kärt Kriisa**. Profile of acylcarnitines, inflammation and oxidative stress in first-episode psychosis before and after antipsychotic treatment. Tartu, 2018, 145 p.
276. **Xuan Dung Ho**. Characterization of the genomic profile of osteosarcoma. Tartu, 2018, 144 p.
277. **Karit Reinson**. New Diagnostic Methods for Early Detection of Inborn Errors of Metabolism in Estonia. Tartu, 2018, 201 p.

278. **Mari-Anne Vals.** Congenital N-glycosylation Disorders in Estonia. Tartu, 2019, 148 p.
279. **Liis Kadastik-Eerme.** Parkinson's disease in Estonia: epidemiology, quality of life, clinical characteristics and pharmacotherapy. Tartu, 2019, 202 p.
280. **Hedi Hunt.** Precision targeting of intraperitoneal tumors with peptide-guided nanocarriers. Tartu, 2019, 179 p.
281. **Rando Porosk.** The role of oxidative stress in Wolfram syndrome 1 and hypothermia. Tartu, 2019, 123 p.
282. **Ene-Ly Jõgeda.** The influence of coinfections and host genetic factor on the susceptibility to HIV infection among people who inject drugs. Tartu, 2019, 126 p.
283. **Kristel Ehala-Aleksejev.** The associations between body composition, obesity and obesity-related health and lifestyle conditions with male reproductive function. Tartu, 2019, 138 p.
284. **Aigar Ottas.** The metabolomic profiling of psoriasis, atopic dermatitis and atherosclerosis. Tartu, 2019, 136 p.
285. **Elmira Gurbanova.** Specific characteristics of tuberculosis in low default, but high multidrug-resistance prison setting. Tartu, 2019, 129 p.
286. **Van Thai Nguyeni.** The first study of the treatment outcomes of patients with cleft lip and palate in Central Vietnam. Tartu, 2019, 144 p.
287. **Maria Yakoreva.** Imprinting Disorders in Estonia. Tartu, 2019, 187 p.
288. **Kadri Rekker.** The putative role of microRNAs in endometriosis pathogenesis and potential in diagnostics. Tartu, 2019, 140 p.